



FLAVONOID CONTENTS FROM SOME FABACEOUS PLANT SPECIES OF RAJASTHAN DESERT

B.B.S. Kapoor* and Veena Purohit

Herbal Research laboratory, P.G. Department of Botany, Dungar College, Bikaner 334001, India
*Corresponding Author Email: bbskapoor@rediffmail.com

ABSTRACT

Evaluation of flavonoid contents from three selected plant species of family Fabaceae like Clitoria ternatea, Sesbania bispinosa and Tephrosia purpurea growing in Rajasthan Desert was carried out. The leaves of selected plants were analysed for flavonoid contents i.e. Quercetin and Kaempferol. Flavonoid contents like Quercetin and Kaempferol were isolated and identified. The maximum total flavonoid contents (1.36mg./gdw) was found in leaves of Clitoria ternatea while minimum (0.91mg./gdw) in leaves of Tephrosia purpurea.

KEY WORDS

Flavonoid contents, Fabaceous plant species, Rajasthan Desert

INTRODUCTION

Rajasthan Desert shows rich phytodiversity.. This region exhibits a great variety of geological, physiographical, climatic, edaphic and biotic conditions and represents diversity of medicinal plant species, which occur on a wide range of habitat. These plant species are good source of phytochemicals of pharmaceutical interest such as flavonoids, sterols, alkaloids, phenolic compounds, sulphides, isothiocyanates, anthocynins, terpenoids etc. These are the active principles which act as antioxidants, anticarcinogenic, antimicrobials and immunity stimulants. A number of plant species have been screened for evaluation of antimicrobial principles like flavonoids [1-7].

MATERIALS AND METHODS

In the present investigation, attempts have been made to investigate the quantitative production of flavonoids in the leaves of Fabaceous plant species like Clitoria ternatea, Sesbania bispinosa and Tephrosia purpurea.

Dried and powdered leaves of the selected plant species were collected from Bikaner district and separately soxhlet extracted with 80% hot ethanol [8]. On a water bath for 24 hrs. Each of the extracts was concentrated and concentrate re-extracted with petroleum ether (Fraction-I), ether (Fraction-II) and ethyl acetate (Fraction-III) in succession. Fraction-III was dried in vacuo and the resultant was hydrolysed with 7% H₂SO₄ for 2 hrs. The mixture was filtered and filterate extracted with ethyl acetate. Concentrated ether and ethyl acetate fraction were applied on TLC plates along with standard reference compounds and the plates developed with the solvent system n-butanol, acetic acid and water (4:1:5) when kaempferol and quercetin were detected.

The compounds were isolated by preparative TLC and crystallized, mp (quercetin $309)^0-311^0$ C and kaempferol 271^0-273^0 C). IR spectra compared well

International Journal of Pharmacy and Biological Sciences (e-ISSN: 2230-7605)



IJPBS | Volume 3 | Issue 3 | JUL-SEPT | 2013 | 209-211

with their authentic samples. Quantitative estimation of flavonoid contents was carried out by method of

Kariyone *et al.* [9] and Naghski *et al.* [10] for quercetin and Mabry *et al.* [11] for kaempferol.

RESULTS AND DISCUSSION

Quercetin and Kaempferol were isolated and identified. Their quantitative estimation is given in the following **Table 1**.

Table 1: Flavonoid contents (mg. /gdw) from leaves of selected plant species

Plants	Quercetin	Kaempferol	Total contents
Clitoria ternatea	0.82	0.54	1.36
Sesbania bispinosa	0.54	0.58	1.12
Tephrosia purpurea	0.39	0.52	0.91

The present investigation shows (**Table 1**) that among all the three samples tested the total flavonoid contents were found to be maximum (1.36mg/gdw) in leaves of *Clitoria ternatea* while minimum (0.91mg/gdw) in *Tephrosia purpurea*.

The maximum quercetin (0.82mg/gdw) in leaves of *Clitoria ternatea* while minimum (0.39mg/gdw) in *Tephrosia purpurea*.

The maximum amount of kaempferol (0.58mg/gdw) in leaves of Sesbania bispinosa while minimum (0.52mg/gdw) in Tephrosia pupurea..

The fabaceous plant species, under study are a potential source of antimicrobial principles. These are resistant to bacterial and fungal attacks due to presence of biologically active substances i.e. flavonoids. These retain potentialities to synthesize the flavonoid contents which are active principles against bacterial as well as fungal pathogens. Due to presence of these secondary products the selected plant species can be used in drug and pharmaceutical industries.

ACKNOWLEDGEMENT

The authors wish to acknowledge the UGC, Bhopal for providing the financial assistance for the project.

REFERENCES

- Nag TN Mathur CS and Goyal SC, Phytochemical studies of *Tribulus terrestris* and *Agave wightii*, contents of primary and secondary products. *Comp. Physiol. Ecol.* 4 157-160 (1979)
- Singh V, Sethia M, Mathur K, Nag TN, Flavonoids of some arid zone plants of Rajasthan. *Ind. J. Pharm. Sci.*, March-April, 88 133 (1988)
- Ahmed-El-Sawi S Abd-El-Megeed HF and Ali AM, Flavonoid and antimicrobial volaties from Adhatoda vasica Nees. Pharmaceuticl and Pharmacological letters 9 (2) 52-56 (1999)
- Ahmed I and Beg Arina Z, Antimicribial and Phytochemical studies on 45 Indian medicinal plants against multi-drug resistant human pathogens. J. of Ethnopharmacology 2 113-123 (2001)
- Kapoor BBS and Ranga P, Flavonoids from Asteraceous medicinal plants of Rajasthan desert. *J. Phytol. Res.* 16(1) 101-102 (2003)



Available Online through

www.ijpbs.com (or) www.ijpbsonline.com

- Kapoor BBS and Kumar S, Herbal plants of Rajasthan desert: A potential source of antimicrobial principles. J. of Aridland studies, Japan 16 425-426 (2005)
- Kapoor, B.B.S. and Mishra, Raksha, Flavonoid Contents from Some Capparidaceous Medicinal Plants of Northwest Rajasthan. *Indian Journal of Pharmaceutical and Biological Research*. 1(2): 9 – 11(2013)
- Subramanian SS and Nagarajan S, Flavanoids of the seeds of *Crotalaria retusa* and *C.striata Curr. Sci.* (India) 38:65 (1969)

IJPBS | Volume 3 | Issue 3 | JUL-SEPT | 2013 | 209-211

- Kariyone T, Hashimato Y. Kinnira M, Microbial studies on plant components.IX.Distribution of flavanoids in plants by paper chromatography. *J. Pharm. Soc.* (Japan), 73 253-256 (1993)
- Naghski J, Fenske (Jr.) CS and Couch IF, Use of paper chromatography for the estimation of quercetin in rutin. J. Pharm. Assoc. 40 613 (1975)
- 11. 11 Mabry TJ, Markham KR. Thomas MB, *The systemic identification of flavonoids*, Springer Verlag, Berlin, 119 (1970).



*Corresponding Author:

B.B.S. Kapoor

Herbal Research laboratory, P.G. Department of Botany, Dungar College, Bikaner 334001, India

*Email: <u>bbskapoor@rediffmail.com</u>