



## EFFECT OF HEAVY SMOKING ON PULMONARY FUNCTION FOR PROFESSIONAL AIR CONDITIONER AND NON- AIR CONDITIONER USERS

R. Lokesh\*<sup>1</sup> and Suganya<sup>2</sup>

\*<sup>1</sup>Assistant professor and <sup>2</sup>internship student  
School of physiotherapy, VISTAS, Chennai

\*Corresponding Author Email: [physiolokesh@gmail.com](mailto:physiolokesh@gmail.com)

### ABSTRACT

**Background:** The cigarette smoking has extensive effect on respiratory function so the risk of respiratory mortality / morbidity is high with smoking. Pulmonary function in young healthy air conditioner (AC) users have 25-75% of vital capacity reduced between the FEF and PEF. **Objectives:** 1. To find out the pulmonary function of professional Air Conditioner users with heavy smoking. 2. To find out the pulmonary function of non-air conditioner with heavy smoking. 3. To compare the pulmonary function for professional Air Conditioner and non-air conditioner users with heavy smoking. **Methodology: setting:** AV modular furniture (India private limit). **study design:** observational study. **Materials & methods:** The pulmonary function tests were assessed on Peak flow meter, Spirometer & Pulse oximeter in 80 subjects (40-AC smokers & 40 - non - AC smokers) and results were compared. **Result:** The mean values for AC smokers PEF - 383.13, Inspiratory capacity - 4562.50, SPO<sub>2</sub> - 95.08 pulmonary function levels were significantly reduced than non-AC smokers. **Conclusion :** The mean PEF, Inspiratory capacity & SPO<sub>2</sub> were lower in AC smokers.

### KEY WORDS

Air Conditioner, smokers, Pulmonary function tests, Peak flow meter, Spirometer & Pulse oximeter.

### INTRODUCTION:

Over 5 million people are killed annually due to smoking. As reported by the 'WHO', by 20th century 100 million people were died and is predicted that by the end of 21st century more than 1 million people will be the victims of smoking<sup>(4)</sup>. It has firm roots over the rural and urban provinces of India in the form of bidis, pipes, cigar etc. <sup>(6)</sup> Consumed mainly in the form of bidis (54%), smokeless tobacco (27%), and cigarettes (9%) <sup>(6)</sup>. The major constituents of cigarette named nicotine can cause arteriosclerosis, coronary artery disease, COPD etc<sup>(11)</sup>. Basic respiratory functions are extensively affected and causes chronic bronchitis, emphysema, bronchial carcinoma <sup>(1,6)</sup>. Pulmonary functions are also damaged due to smoking and lead to oxygen insufficiency in cells as a result of elevated carbon

monoxide in the blood and decrease the oxygen carrying capacity of hemoglobin <sup>(12)</sup>.

Considerable evidence has been obtained that the neutrophils are important in mediating lung damage due to smoking. It has been observed that the circulating leucocyte counts and broncho alveolar lavage(BAL) neutrophil counts are much higher in the current smokers as compared to the ex-smokers/non-smokers and there is an inverse correlation between circulating leucocyte count and one second forced expiratory volume(FEV1) and forced vital capacity(FVC). The experimental studies carried out in hamsters came to a conclusion that even four hours of exposure to cigarette smoke causes extensive recruitment of leucocytes in to the airway walls. In

healthy smokers the increased in vitro elastase activity has been seen in cell free BAL fluid<sup>(18)</sup>.

AC is a device that is using to provide cooling indoor by condensation the atmospheric water vapour and thus reducing the humidity of air<sup>(9)</sup>. AC's are used extensively these days Indoor as well as while travelling. Over expose / inhalation of cold dry air an ultimately lead to alteration in the pulmonary function. Increased prevalence of IgE induce sensitization and hyper sensitivity pneumonitis are reported in the person exposed to aerosols of contaminated AC<sup>(10)</sup>. Mucous membrane irritation, breathing difficulties, irritated skin and constitutional or neurological symptoms such as head ache and fatigue are complaints faced by most of the AC users<sup>(9)</sup>.

Various studies conducted by a team of doctors from Indian Medical Colleges has come to a conclusion that people, who are over exposed to air-conditioned spaces are more likely to have respiratory ailments like coughing, wheezing and breathlessness, and the frequency of such disease are high among them. Surveys conducted by several international and Indian agencies have brought out the ill effects of excessive use of AC on humans.

Studies were conducted by 'Mendel' with the US department of energy's Lawrence Berkeley National Laboratory. On the health effects of air conditioning systems. They came to a conclusion that worsening asthma problems and allergies are the two-health issue that can stem from contaminated AC units.

---

#### **METHODOLOGY:**

---

- Sample size: 80
- Study location: AV modular furniture (India private limit )
- Study duration: single time study
- Study design: observational study
- Study sampling: purposive sampling
- Study type: comparative study

#### **INCLUSION CRITERIA:**

- Age: 20 to 30 years
- Gender: male only
- Condition: AC and Non- AC smokers
- Number of smoking: More than 200(cigarettes /bidis) per year.

#### **EXCLUSION CRITERIA:**

- History of acute or chronic respiratory disorders
- History of Cardiac disease

- The patient who all are undergoing medication for any cardiac or respiratory problem
- Regularly exercising individual
- Chronic alcoholic.

#### **TOOLS:**

##### **PULSEOXIMETER:**

A Pulse oximeter is a non-invasive device that estimates the amount of oxygen in blood. It does so by sending infrared light into capillaries in fingers, toe or earlobe.

##### **PEAK FLOW METER**

A peak flow meter is an inexpensive portable, hand held device, it used to measure maximum speed of expiration.

##### **SPIROMETER:**

A spirometer is an apparatus for measuring the volume of air inspired and expired by the lungs.

#### **OUTCOME MEASURES**

- Oxygen saturation level
- Peak expiratory flow rate
- Inspiratory capacity

#### **PROCEDURE:**

Totally 100 subjects were screened by means of questionnaire. Out of which 80 subjects were included in the study according to inclusion and exclusion criteria. After getting informed consent, they were divided into two groups according their experience.

First the personal data will be taken with include age, gender, history of smoking, weight and height. The participants were divided into two groups. Group A and Group B each consisting of 40subjects. Group A – smoker who did not use AC, Group B – smoker who had used AC (air conditioner. Both groups were assessed of their respiratory parameters, they are included Peak Expiratory Flow Rate (PEFR), inspiratory capacity and saturation level and data were collected, tabulated and analyzed statistically for evaluating the respiratory function.

#### **PEAKEXPIRATORY FLOW RATE:**

The subjects were asked to sit comfortably on chair(erect position).The complete procedure was explained the subjects were instructed to breath in fully and deeply inspiration with nostrils closed,seal the lips around the sterile mouth piece of peak flow meter and forcefully expire the air out as fast and as far as possible(they asked to do 2-3 times repeatedly) and then the maximum level will be recorded.

**INSPIRATORY VITAL CAPACITY:**

The subjects were asked to sit comfortably on chair (erect position). The complete procedure was explained. The subjects were asked to inspire through the mouthpiece as much as possible (they asked to do 2-3 times repeatedly) and then the maximum reading was recorded.

**OXYGEN SATURATION LEVEL:**

The subjects were asked to sit in a comfortable position (erect position). A clip like device will be placed on finger. Within 5 to 10 seconds the oxygen saturation level will be monitored and then reading will be recorded.

**STATISTICAL ANALYSIS:**

Statistical analysis was done by using the single t test. The significance was drawn at a P (probability) value of 0.05. The outcome of pulmonary function tests were presented as a mean  $\pm$  standard error of difference for each of the parameter. The two groups were compared by applying single t test and P value of less than 0.05 (\*P < 0.05) was considered as significant.

T test is often called student's t test in the name of its founder "student". T test is used to compare two different sets of values. It is generally performed on a

small set of data. T test is generally applied to two normal distributions which have a small set of values. This test compares the mean of two samples. T test uses means and standard deviation of two samples to make a comparison. The formula for T test is given below:

$$t = \frac{\bar{x}_1 - \bar{x}_2}{\sqrt{s^2 \left( \frac{1}{n_1} + \frac{1}{n_2} \right)}}$$

Where,

x1 = Mean of first set of values

x2 = Mean of second set of values

S1 = Standard deviation of first set of values

S2 = Standard deviation of second set of values

n1 = Total number of values in first set

n2 = Total number of values in second set


The standard deviation of a sample is known as S and is calculated using:

$$s = \sqrt{\frac{\sum (x - \bar{x})^2}{n - 1}}$$

P value < 0.05 was taken as a level of significance.

**TABLE 1: GROUP A  
SMOKING WITH NON-AC USERS (GROUP A)**


LUNG PARAMETERS	MEAN	SD	T	P
PEFR	434.00	59.82	45.8851	0.0001
INSPIRATORY RATE	4620.00	443.88	65.8280	0.0001
SPO2	96.45	4.70	129.9094	0.0001



**GRAPH-1 SHOWS PEFR, INSPIRATORY CAPACITY, SPO<sub>2</sub> LEVEL OF SMOKING WITH NON-AC USERS (GROUP A)**

**TABLE 2: GROUP B  
SMOKING WITH AC USERS (GROUP B)**


LUNG PARAMETERS	MEAN	SD	T	P
PEFR	383.13	83.27	29.0986	0.0001
INSPIRATORY	4562.50	642.39	44.9197	0.0001
SPO2	95.08	6.60	91.0553	0.0001



**GRAPH-2 SHOWS PEFR, INSPIRATORY CAPACITY, SPO<sub>2</sub> LEVEL OF SMOKING WITH AC USERS (GROUP B)**

**TABLE 3: COMPARISON BETWEEN GROUP A AND B**

LUNG PARAMETERS	SMOKERS WITH NON-AC USERS (GROUP A) MEAN VALUE	SMOKERS WITH AC USERS (GROUP B) MEAN VALUE
PEFR	434.00	383.13
INSPIRATORY	4620.00	4562.00
SPO2	96.45	95.08



**GRAPH-3 SHOWS PEFR, INSPIRATORY CAPACITY, SPO<sub>2</sub> LEVEL OF BOTH NON-AC SMOKERS AND AC SMOKERS**

#### RESULT:

Table 1 shows that Group A (Non-AC smokers) mean value  $\pm$  standard error of difference of PEFR (434.00 $\pm$ 9.46), inspiratory rate (4620.00 $\pm$ 70.18) and SPO<sub>2</sub> (96.45 $\pm$ 0.74) is significant. Table 2 shows that Group B (AC smokers) mean value  $\pm$  standard error of difference of PEFR (383.13 $\pm$ 13.17), inspiratory capacity (4562.50 $\pm$ 101.57), SPO<sub>2</sub> (95.08 $\pm$ 1.04) is significant. AC smokers mean value (PEFR-383.13, Inspiratory capacity-4562.50, SPO<sub>2</sub>-1.04) than Non-AC smokers (PEFR-434.00, Inspiratory capacity-4620.00, SPO<sub>2</sub>-96.45)

#### DISCUSSION:

This study is focused to see the changes occur between effect of heavy smoking on pulmonary function for professional Air Conditioner and non-air conditioner users. The result of the current study the mean values of the spirometry variables were found to be higher in non AC smokers than AC smokers and also the difference in PEFR was significantly higher in the non AC smokers.

Drawback of this study is that the fall in PEFR indicates obstructive lung changes and fall in FVC indicates restrictive lung changes. smoking and intensive use of AC "s appeared to be positively related to atopic sensitization and enhanced eosinophil activity. As we see the P-value of group A and group B in different pulmonary function test.

Hence group A which shows effective significance for all the tests. The way of test and the comfort which gave them a better improvement and made changes in their

professional life. Study by keele CA et al, PEFR depends on the expiratory efforts exerted during forceful expiration as well as status of airway, And it reflects mainly the caliber of the bronchi and larger bronchioles . which are subjected to reflex broncho constrictions.

#### RECOMMENDATION:

1. Sample size can be increased.
2. Using of more parameters should be useful in future study like humidity level, culture swap from AC to know the growth of bacteria and fungi .
3. Female can also be included.
4. AC users might do pulmonary function tests regularly to detect any changes at an earlier stage.
5. Use advance techniques like computerized spirometer.
6. Age groups can be changed.

#### LIMITATION:

1. Less sample sizes
2. Less parameters used
3. Only males included

#### CONCLUSION:

Smoking in any form, bidi/cigarette affects the pulmonary function. Pulmonary function plays a vital role for the survival of human being. If pulmonary function is affected oxygen demand will be increased. It may lead to several health issues.

This study concludes, pulmonary function parameters like PEFR, Inspiratory capacity, SPO<sub>2</sub> levels were markedly reduced for both groups. But comparatively

Group B (ie) AC smokers pulmonary functions was much reduced than non-AC smokers.

---

**REFERENCE:**

1. HARITA P VYAS, RUTVEE P VINCHHI, MEGHA S SHETH, NEETA J VYAS, Comparison of pulmonary function among smokers and non- smokers - a retrospective study, International journal of medical science and public health[IJMSPH], Volume 3, Issue 10,2014.
2. FARAH KHALIQ, SAMEER SHARMA AND O.P. TANDON, Pulmonary function in Air conditioner users, Indian journal physio protocol [2006]
3. CHANDRASHEKHAR D DAMAGE, JAYANT DEORE, ANITA J JADHAV, Pulmonary function test in professional Air conditioner users, IOSR journal of dental and medical science, volume 15, issue 4,[april 2016].
4. RUBEENA BANO, MAHAGAONKAR AM, KULKARNI NB, NADEEM AHMAD, NIGHUTE S, Study of pulmonary function tests among smokers and nonsmokers in a rural area, Paravara medical review 2009
5. HANI A. NAWAFLEH, SHALABIA A1- STAYED ABO ZEAD, DUA'A FAYEZ A1- MAGHAIREH, Pulmonary function test: the value among smokers and nonsmokers, Health science journal, volume 6, issue 4, [oct - Dec 2012].
6. RUBEENA BANU, NADEEM AHMAD, AM MAHAGAONKA, study of pulmonary function in smokers and non - smokers in sugarcane harvesters in rural Maharashtra, Walawalkar international medical journal, volume 1, issue 1, [2014].
7. LAXMIKANT J BORSE, RASIKA D YADAV, HITESH K MODAK, LALITA M CHANDAN, Pulmonary function test in young healthy male exposed to air-conditioned work environment, International journal of health sciences and research, volume 2, issue 6, September 2012.
8. R. BABITHA, R. RANGARAJAN, M. MUHIL, M.G BARAVARA JAIAH, Pulmonary function tests in air conditioner users, Journal of clinical and diagnostic research, volume 5, june 2011.
9. VIDYA G, KUMAR B.A, KALPANA M, CHAND K, Pulmonary function tests in air conditioner users, International journal of medicine and bio medical research. volume 3, issue 2, may - august 2014.
10. MOHAMMED JEELANI, MOHAMMAD AHMED, Effect of air conditioner on pulmonary functions in healthy males in and around raichur city, Journal of evidence-based medicine and health care. Volume 2, issue 19, May 2015.
11. KYOCHUL SEO, PhD, SEUNG HWAN PARK, PhD, KWANGYONG PARK, MS, Effects of diaphragm respiration exercise on pulmonary function of male smokers in their twenties, Journal physical therapy science 2015.
12. HYUN - JU JUN, PT, PhD, KI - JONG KIM, PT, PhD, KI - WON NAM, PT, PhD, CHANG – HEON KIM, PT, PhD, Effects of breathing exercises on lung capacity and muscle activities of elderly smokers, Journal of physical therapy science , 2016.
13. Dr. R MANIMOZHI MALATHI, Dr. D RAJKUMAR, Dr.E GNANA DESIGAN, Dr. K BALUMAHENDRAN, A study of pulmonary function tests among smokers in rural areas, IOSR journal of dental and medical sciences , volume 16, issue 12, dec. 2017.
14. AGRAWAL PB, SARODE VS, SHAH SH, KOWALE AN, A comparative study of pulmonary function tests in air - conditioner users and non-users, Scholars journal of applied medical sciences, 2017.
15. SABADE SB, B.B. VIKHE\*, BORADE NG\*\*, Pulmonary ventilation in air conditioner users in paravara rural hospital, Paravara medical review [2013].
16. ANA PAULA COELHO FIGUEIRA FREIRE, DIONEI RAMOS, The role of professional physical therapist in smoking cessation: A literature update, General medicine [los angeles] an open access journal, ISSN , Volume 5, issue 2, 2017.
17. JOSHUA SILL MD\*, Tobacco cessation and prevention, Chest net organization [April 2016].
18. C R K HIND, G A TENNENT, H JOVCE, M B PEPYS, N B PRIDE, Plasma leucocyte elastase concentration in smokers, J Clin Pathol [1991].
19. BRENDA W, CAMPBELL JENKINS, DANIEL F. SARPONG, CLIFTON ADDISON, MONIGUE S. WHITE, DEMARE A. HICKSON, WENDY WHITE and CECIL BURCHFIEL, Joint effects of smoking and sedentary life style on lung function in african americans: the jackson heart study cohort, International journal of environmental research and public health . [2014].
20. S.P. CHOUDHARI, R.S. DOIPHODE, U. S. ZINGADE, Study of pulmonary functions in air-conditioned car drivers: A cross sectional study, IOSR journal of dental and medical science. [jan-2014].
21. MANIKANDAN S, ANANDHALAKSHMI S, NAGESWARI AD, Comparison of the effects of various modes of smoking on the pulmonary functions in healthy volunteers, Asian journal of pharmaceutical and clinical research. volume 8, issue 3,2015.
22. J.L. WRIGHT, J. HOBSON, B.R. WIGGS, Effect of cigarette smoking on structure of the small airways, Lung [1987].
23. MATURI LAVANYA, RAJANAND GAIKWAD, The effect of air conditioner on pulmonary functions in young adults, International journal of integrative medical sciences, volume 4(6):501-06, 2017.
24. SANDIP M. HULKE, AVINASH THAKARE, pulmonary functions in air conditioner users, Medical journal of Dr. D. Y. Patil university, volume 6, issue -1, 2013.
25. ANU T. E.<sup>1\*</sup>, BIJU BAHULEYAN<sup>2</sup>, Analysis of dynamic pulmonary functions in air-conditioned work environment, International journal of research in medical sciences, 2016 july;4(7).



26. RAFAEL LANIADO - LABORIN, Smoking and chronic obstructive pulmonary disease (COPD). Parallel epidemics of the 21<sup>st</sup> century, International journal of environmental research and public health, 2009.
27. ROBERT J. HANCOX<sup>1</sup>, ANDREW R. GRAY, The effect of cigarette smoking on lung function in young adults with asthma, American journal of respiratory and critical care medicine, volume -194 number 3, 2016 august.
28. RACHEL E JORDAN, KAR KEUNG CHENG, Passive smoking and chronic obstructive pulmonary disease; cross sectional analysis of data from the health survey for england, BMJ Open 2011
29. LEAH JAYES, BSc; PATRICIA L. HASLAM, PhD, SmokeHaz systemic reviews and meta-analysis of the effects of smoking on respiratory health, chest 2016 july .
30. PRUTHI, N., & MULTANI<sup>2</sup>, N. K, Influence of age on lung function tests, Journal of exercise science and physiotherapy, volume -8, no. 1: 1-6, 2012.

**\*Corresponding Author:**

**R.Lokesh\***

Email: [physiolokesh@gmail.com](mailto:physiolokesh@gmail.com)