



## PROBIOTIC HOLD PROMISES FOR GLOWING AND HEALTHY SKIN

Verinder Wahla and Garima Verma\*

Kanya Gurukul campus Haridwar- 249407, India.

\*Corresponding Author Email: [garimav467@gmail.com](mailto:garimav467@gmail.com)

### ABSTRACT

*Probiotic bacteria are used to treat or prevent broad range of skin disease, condition and syndrome. In addition, there are area of medical use that have been proposed for future probiotic application in skin diseases this review have provide evidence of probiotic effectiveness for the treatment and prevention of skin aging, acne, skin allergy and atopic dermatitis that cure with the help of regular intake of probiotic and dietary food products or the probiotic bacteria that are active against many skin diseases.*

### KEY WORDS

*Atopic dermatitis, probiotics, dietary supplements, S. epidermidis.*

### INTRODUCTION

Probiotics are “live microorganisms, which, when administered in adequate amounts, confer a health benefit to the host” (Anon, 2006). It has been speculated that the skin status could benefit from reinforced gut homeostasis (Salminen et al., 2005). Diet variations determine individual characteristics of intestinal microflora, according to age, feeding, lifestyle, interactions among numerous constituents of the same flora and pathologic conditions. Probiotic formulations are becoming increasingly available for healthy skin care, prevention and treatment of skin diseases, and anti-aging benefits thus representing an emerging area for skin health (Cinque et al., 2011). The advantages of probiotic treatment (probiotic-prevention or probiotic therapy) are that this method is efficient to the patients and also it has no side effect on them. Therefore, probiotic-therapy for the skin is potentially comparable to the ordinary methods of treatments.

In modern lifestyle negatively influences the intestinal ecosystem, and there may be cumulative degradation of the intestinal microflora. Linear changes in environmental conditions and lifestyle may lead to nonlinear changes in the gut flora possibly to an increasing susceptibility to skin diseases (Schmidt,

2004). A number of clinical studies suggest that probiotic strategies induce systemic effects that extend beyond the gut and may even affect selected functions of the skin (Ouweland et al., 2002). Many experimental studies have shown that probiotics exert specific influences in the intestinal on epithelial cells and immune cells with antiallergic potential (Caramia et al., 2008). Also, an emerging approach in dermatology to help preventing and treating skin conditions, including the external signs of aging, acne, rosacea, yeast and bacterial infections, psoriasis, and dermatitis, is represented by topical probiotics, as shown by the growing market place for topical probiotic formulations available for skin care and antiaging benefits (Cinque et al., 2011). Totally, probiotics exert their health effects to the skin directly through cutaneous formulations or indirectly through dietary supplementary formulations. This review is related to the beneficial effect of probiotic bacteria on skin aging, atopic dermatitis, acne and skin allergy.

## 1. SKIN AGING



**Figure1: wrinkles on skin (Source: <https://www.yahoo.co.in>)**

Aging skin involve a complex interplay between intrinsic aging including genetic and hormonal influence and extrinsic aging influence enviornmental factor such as uv light, pollution and cigarette smoking. UV radiation is considered the strongest precipitator of extrinsic aging. As public knowledge has increase dregarding the link between sun exposure and photoaging, interest in preventing and treating the adverse effects of UV radiation has escalated. Early studies suggest that probiotics and their metabolites might alter several aspects of skin aging and also help to maintain skin homeostasis by protecting against UV-induced skin damage (J. h. jeong *et al.*, 2016). Healthy, normal skin exhibits a slightly acidic pH in the range of 4.2- 5.6, which aids in the prevention of pathogenic bacteria colonization, regulation of enzyme activity, and maintenance of a moisture-rich environment (Mauro, 2006). However, after the age of 70, the pH of skin raises significantly, stimulating protease activity (Hachem *et al.*, 2003). Probiotic metabolism frequently produces acidic molecules, lowering the pH of the surrounding environment (Cinque *et al.*, 2010) as seen with *Lactobacilli* producing free fatty acids (FFAs) and conjugated linoleic acid (CLA) during the fermentation process. Therefore, the use of probiotics may work to restore the normal skin pH and consequently return protease activity levels closer to those seen in young and healthy skin.

## 2. ATOPIC DERMATITIS



**Figure 2: Atopic dermatitis (Source: <https://www.yahoo.co.in>)**

Atopic dermatitis is the most common chronic skin condition in infants and children, with a prevalence of 10–20% (Kliegman *et al.*, 2015). The term atopy refers to a genetic predisposition to become sensitized and to mount an IgE response to allergens. Atopic dermititis has been linked to food hypersensitivity, especially milk and egg protein. It is also defined as “a skin condition characterized by intense dryness of the skin, pruritus, and chronic erythematous lesions with a relapsing course” (Del Giudice jr *et al.*, 2002). Atopic dermatitis is a chronic relapsing inflammatory skin disease that usually starts during the first years of life. In patients with the disease, the quality of skin is severely affected, and this is closely linked to a reduced quality of life. AD has three stages, non-atopic dermatitis, “True” atopic dermatitis and autoallergic atopic dermatitis (Bieber, 2008). Non-atopic dermatitis is a non-IgE associated form of AD. However, it can transform into “true” atopic dermatitis due to the development of sensitization of the skin to IgE. “True” atopic dermatitis can develop into autoallergic atopic dermatitis by scratching. Scratching damages the skin cells which releases autoantigens. These autoantigens induce IgE auto antibodies which make the inflammatory response worse (Altrichter *et al.*, 2008). A probiotic bacteria *S. epidermidis* can be used to improve atopic dermatitis by combatting *S. aureus* infection. *S. aureus* is found in 90% of atopic dermatitis lesions and worsens the symptoms of atopic dermatitis. Esp secreted by *S. epidermidis* can inhibit *S. aureus* and potentiate the effects of skin antimicrobial peptide produced by the host.

### 3. ACNE



**Figure 3: Acne (Source: <https://www.yahoo.co.in>)**

Acne (vulgaris) is a common disease that mainly affects teenagers. About 80% of the people acquire acne in their adolescence (Cunliffe and Gollnick, 2001). People can experience acne symptoms for decades, nearly 50% of these people suffer from acne in their 20s (Al Robaee, 2005 and Galobardes *et al.*, 2005). Acne vulgaris is multifactorial condition and is characterized by hyper colonization with *P. acnes* inflammation and immune responses. Propionibacteria have been shown to have adjuvant and antitumor activities (Ingham *et al.*, 1992) and also same species are of pathogenic relevance in acne and folliculitis (Leyden *et al.*, 1998). The symbiotic ability of probiotic bacteria and Konjac glucomannan hydrolysates to inhibit the growth of *P. acnes* in an in vivo study has been recently reported suggesting that the development of a new alternative involving probiotic therapy for reducing acne episodes in vivo could be encouraging (Al-ghazzawi and Tester., 2009). The local burden of lipid peroxidation in acne is high, such that it appears to place a great demand upon blood-derived antioxidants (Bowe and Logan, 2010). Some recent studies have shown that orally consumed prebiotic and probiotics can reduce systemic markers of inflammation and oxidative stress (Mikelsaar and Zilmer., 2009). Then the ability of oral probiotics to limit systemic oxidative stress may be an important therapeutic pathway (Fu *et al.*, 2010). Oral administration of probiotics regulates the release of inflammatory cytokines within the skin (Hacini-Rachinel *et al.*, 2009), and a specific reduction in interleukin-1 $\alpha$  (Cazzola *et al.*, 2010) would be of potential benefit in acne (Bowe and Logan, 2011). Probiotics have been shown to directly inhibit *P. acnes* through the production of antibacterial proteins.

### 4. SKIN ALLERGY



**Figure 4: Allergic diseases (Source: <https://www.yahoo.co.in>)**

The role of probiotics in prevention of allergic disease is still not clearly established. It has been shown that probiotics exert specific anti-allergic effects on epithelial cells and immune cells. The allergen penetration through the skin can lead to a systemic sensitization involving the intestinal mucosa. Also, it has been reported that inflammation plays an important role in photo ageing of human skin in vivo (Pillai *et al.*, 2005). Intervention strategies have been elaborated to balance the intestinal microecology with oral administration of probiotics (Ogden and Bielory., 2005). The composition of the intestinal microflora is the main element in allergic diseases, and species of probiotics could play a role in the development of healthy immunity response, participating in enteric microecology, preventing, and potentially treating allergic diseases (Zutavern *et al.*, 2006). Dietary supplementation using particular probiotic strains has been found to increase interferon activity in the blood of human, and children born to families who consume traditional *Lactobacillus* rich fermented foods experience fewer allergies than those from families who consume more sterile foods (Caramia *et al.*, 2008). It has been confirmed that children who develop allergies present low levels of *bifidobacteria*, gram-positive aerobic organisms and enterococci, but more elevated levels of clostridia and *S. aureus* in their enteric microflora. Increased fecal levels of i-caproic acid that are indicators of high level of *Clostridium difficile*, suggest that the enteric microflora could be changed in allergic children. Many factors exert the alteration of gastrointestinal flora equilibrium such as genetic factors, diet, and infection frequency, antibiotic therapies, passive smoke, pollution, vaccinations, psychologic stresses, reduction of immunologic activity,

and development of allergic diseases (Halken, 2004). For example, antibiotic therapy leads to changes in intestinal, bacterial, and fungal components and also to over growth of *Candida albicans* which can secrete potent prostaglandin-like immune response modulators and subsequently promote the development of allergic events in distal mucosal sites such as the skin (Noverr *et al.*, 2005). It should be noted that not all probiotic microorganisms have the same immunologic properties and specific probiotic microorganisms of the developing intestinal microflora that most influence skin health need to be defined because specific deviations in intestinal microflora may exert predisposition to allergic disease (Caramia *et al.*, 2008). Totally, allergic responses may also be affected by dose and viability of the probiotic and these factors in turn potentially modified by the host environment (Wickens *et al.*, 2008).

## CONCLUSION

The role of probiotic bacteria in the prevention and treatment of skin diseases such as atopic dermatitis, acne, skin aging and allergic reactions. However, probiotic bacteria are unanimous but more evidences are needed to prove the impact of probiotic in the treatment of skin disease especially for allergic disorders. Probiotic bacteria *Lactobacillus*, *S. epidemis*, *Bifidiobacteria* provide protective shield against skin diseases. Oral administration of probiotic also plays significant role against bacteria which cause skin disease like acne and skin infection. Regular intake of probiotic and dietary food products can reduce minimum risk of skin related diseases last but not least further investigations are also required, in vitro and in clinical trials, with different species and strains of probiotic microorganisms in relevance to their effects on the skin.

## REFERENCES

1. Al Robaee, A. A. Prevalence, knowledge, beliefs and psychosocial impact of acne in University students in Central Saudi Arabia. Saudi Medical Journal, 26(12), 1958-1961, (2005).
2. Al-Ghazzewi, F. H. and Tester, R. F. Effect of konjac glucomannan hydrolysates and probiotics on the growth of the skin bacterium *Propionibacterium acnes* in vitro. International Journal of Cosmetic Science. 32:139-142, (2009).
3. Altrichter, S., Kriehuber, E., Moser, J., Valenta, R., Kopp, T., & Stingl, G. Serum IgE autoantibodies target keratinocytes in patients with atopic dermatitis. Journal of Investigative Dermatology, 128(9), 2232-2239, (2008).
4. Anon. Guidelines for the evaluation of probiotics in food. Report of a Joint FAO/WHO Working Group on Drafting Guidelines for the Evaluation of Probiotics in Food, (2006).
5. Bieber, T. Atopic dermatitis. The New England Journal of Medicine, 358, 1483-1494, (2008).
6. Bowe, W. P., & Logan, A. C. Clinical implications of lipid peroxidation in acne vulgaris: old wine in new bottles. Lipids Health Dis, 9, 141, (2010).
7. Caramia, G., Atzei, A. and Fanos, V. Probiotics and the skin. Clinical Dermatology, 26:4-11, (2008).
8. Caramia, G., Atzei, A. and Fanos, V. Probiotics and the skin. Clinical Dermatology, 26:4-11, (2008).
9. Cazzola, M., Tompkins, T. A. and Matera, M. G. Immunomodulatory impact of a synbiotic in T(h)1 and T(h)2 models of infection. Therapeutic Advance in Respiratory Disease. 4:259-270, (2010).
10. Cinque, B., Palumbo, P., La Torre, C., Melchiorre, E., Corridoni, D., Miconi, G., Di Marzio, L., Cifone, M.G. and Giuliani, M. Probiotics in aging skin. In: Text book of Aging skin, Farage, M.A., Miller, K.W., and Maibach, H. I., eds. Springer, Berlin, (2010).
11. Cinque, B., Torre, C., Melchiorre, E., Marchesani, G., Zoccali, G., Palumbo, P., Marzio, L. D., Masci, A., Mosca, L., Mastromarino, P., Giuliani, M. and Cifone, M. G. Use of probiotics for dermal applications. In: Probiotics, microbiology monographs. Liong, M.-T., Ed., Springer Verlag, Berlin, Heidelberg, (2011).
12. Cunliffe, W. J., & Gollnick, H. P. Acne: Diagnosis and management. Taylor & Francis, (2001).
13. del Giudice Jr, M. M., De Luca, M. G. and Capristo, C. Probiotics and atopic dermatitis. A new strategy in atopic dermatitis. Digestive and Liver Disease, 34:S68-S71, (2002).
14. Fu, Y. R., Yi, Z. J., Pei, J. L. and Guan, S. Effects of *Bifidobacterium bifidum* on adaptive immune senescence in aging mice. Microbiology. Immunology. 54:578-583, (2010).
15. Galobardes, B., Davey Smith, G., Jeffreys, M., and McCarron, P. Has acne increased? Prevalence of acne history among university students between 1948 and 1968. The Glasgow Alumni Cohort Study. British Journal of Dermatology, 152(4), 824-825, (2005).
16. Gollnick, H.O., Cunliffe, W., Berson, D., Dreno, B., Finlay, A., Leyden, J. J., Shalita, A. R., & Thiboutot, D. Management of acne: a report from a global alliance to improve outcomes in acne. Journal of the American Academy of Dermatology, 49(1), S1-S37, (2003).
17. Hachem, J. P., Crumrine, D., Fluhr, J., Brown, B. E., Feingold, K. R., & Elias, P. M. pH directly regulates epidermal permeability barrier homeostasis, and

- stratum corneum integrity/cohesion. *Journal of Investigative Dermatology*, 121(2), 345-353, (2003).
18. Hacini-Rachinel, F., Gheit, H., Le Ludec, J. B., Dif, F., Nancey, S., & Kaiserlian, D. Oral probiotic control skin inflammation by acting on both effector and regulatory T cells. *Plus, One*, 4(3), e4903, (2009).
  19. Halken, S. Prevention of allergic disease in childhood: Clinical and epidemiological aspects of primary and secondary allergy prevention. *Pediatric Allergy and Immunology*, 4–5:9–32, (2004).
  20. Ingham, E., Eady, E. A., Goodwin, C. E., Cove, J. H., & Cunliffe, W. J. Pro-Inflammatory Levels of Interleukin-1 $\alpha$ ; -Like Bioactivity Are Present in the Majority of Open Comedones in Acne Vulgaris. *Journal of investigative dermatology*, 98(6), 895-901, (1992).
  21. Ji Hye Jeong, Chang Y. Lee & Dae Kyun Chung. Probiotic lactic acid bacteria and skin health. *Critical Review in Food Science and Nutrition*. 56:2331-2337, (2016).
  22. Kliegman, R. M., Behrman, R. E., Jenson, H. B. and Stanton, B. F. *Nelson Textbook of Pediatrics*. Saunders, Philadelphia, (2015).
  23. Leyden, J.J., McGinley, K.J. and Vowels, B. *Propionibacterium acnes* colonization in acne and nonacne. *Dermatology*, 196(1), 55-58, (1998).
  24. Mauro, T. SCpH: Measurement, origins and functions. In: *Skin Barrier*. Elias, P. and Feingold, K., Eds., Taylor & Francis, New York, (2006).
  25. Mikelsaar, M., & Zilmer, M. *Lactobacillus fermentum* ME-3-an antimicrobial and antioxidative probiotic. *Microbial ecology in health and disease*, 21(1), 1-27, (2009).
  26. Noverr, M. C., Falkowski, N. and McDonald, R. A. Development of allergic airway disease in mice following antibiotic therapy and fungal microbiota increase: Role of host genetics, antigen, and interleukin-13. *Infection and Immunology Journal* 73:30–38, (2005).
  27. Ogden, N. S. and Bielory, L. Probiotics: A complementary approach in the treatment and prevention of pediatric atopic disease. *Current Opinion in Allergy and Clinical Immunology*, 5:179–184, (2005).
  28. Ouwehand, A.C., Salminen, S. and Isolauri, E. Probiotics: An overview of beneficial effects. *Antonie van Leeuwenhoek*, 82:279–289, (2002).
  29. Pillai, S., Oresajo, C. and Hayward, J. Ultraviolet radiation and skin aging: Role of reactive oxygen species, inflammation and protease activation, and strategies for prevention of inflammation induced: A review. *International Journal of Cosmetic Science*, 27:17–34, (2005).
  30. Salminen, S. J., Gueimonde, M. and Isolauri, E. Probiotics that modify disease risk. *The Journal of Nutrition*, 135:1294–1298, (2005).
  31. Schmidt, W. P. Model of the epidemic of childhood atopy. *Medical Science Monitor*, 10:5–9, (2004).
  32. Wickens, K., Black, P. N., Stanley, T. V., Mitchell, E., Fitzharris, P., Tannock, G. W., Purdie, G. and Crane, J. A differential effect of probiotics in the prevention of eczema and atopy: A double-blind, randomized, placebo-controlled trial. *Journal of Allergy and Clinical Immunology*. 122:788–794, (2008).
  33. Zutavern, A., Brockow, I. and Schaaf, B. Timing of solid food introduction in relation to atopic dermatitis and atopic sensitization: Results from a prospective birth cohort study. *LISA Study Group. Pediatrics* 117: 401–411, (2006).

Received:04.05.18, Accepted: 07.06.18, Published:01.07.2018

**\*Corresponding Author:**

**Garima Verma\***

Email: [garimav467@gmail.com](mailto:garimav467@gmail.com)