



Probiotics Against Food-Borne Bacterial Pathogens: A Review and Future Directions

Mary Sandeepa Gujjula*

Department of Biotechnology, Vikrama Simhapuri University, Nellore-524 320, Andhra Pradesh, India

Received: 14 Oct 2021 / Accepted: 9 Nov 2021/ Published online: 01 Jan 2022

*Corresponding Author Email: deepavsu2013@gmail.com

Abstract

Food-borne bacterial pathogens represent a significant public health concern worldwide, with their prevalence exacerbated by antibiotic resistance. As traditional treatment options become less effective, alternative strategies are urgently needed. Probiotics, live microorganisms conferring health benefits to the host, have emerged as promising candidates for combating these pathogens. This comprehensive review examines the development of probiotics as a preventive and therapeutic approach against food-borne bacterial pathogens. We delve into the mechanisms through which probiotics exert antimicrobial effects, including competitive exclusion, antimicrobial substance production, and host immune modulation. Additionally, we evaluate the efficacy of various probiotic strains against common food-borne pathogens such as *Salmonella*, *Escherichia coli*, *Listeria monocytogenes*, and *Campylobacter jejuni*. Furthermore, we address the challenges associated with probiotic development and application in food safety, including strain selection, viability, stability, and regulatory considerations. Finally, we discuss future research directions and the potential integration of probiotics into food production and processing practices. This review provides valuable insights into the current state and future prospects of probiotics in mitigating the risks posed by food-borne bacterial pathogens.

Keywords

Probiotics, food-borne pathogens, bacteria, MRSA.

1. INTRODUCTION

Food-borne illnesses remain a significant global health burden, with bacterial pathogens being a leading cause of outbreaks. Traditional antimicrobial therapies are facing challenges due to the emergence of antibiotic-resistant strains, necessitating the exploration of alternative approaches. Probiotics, defined as live microorganisms that confer health benefits to the host when administered in adequate amounts, have garnered attention for their potential in food safety and public health. The development of probiotics against foodborne bacterial pathogens is a critical area of research aimed at enhancing food safety and reducing the risk of foodborne illnesses. Probiotics, such as *Lactobacillus* and *Bifidobacterium* species, have been extensively studied for their ability to combat foodborne pathogens like *Escherichia coli* and methicillin-resistant *Staphylococcus aureus* (MRSA) [1].

These probiotics have shown significant antimicrobial activity against various pathogens, thereby offering a promising approach to mitigating the spread of harmful bacteria through food. Additionally, probiotics have been found to play a crucial role in modulating the immune system, maintaining the mucosal barrier, and preventing conditions like food allergies and diarrhoea.

Research has also focused on incorporating probiotics into food packaging materials to control foodborne microorganisms and improve food safety [2]. By encapsulating probiotics in biopolymer matrices, active food packaging materials have been developed as an alternative to traditional packaging, extending the shelf life of food products and promoting health benefits. Furthermore, the use of probiotics in both dairy and non-dairy products has been explored to meet consumer

demands and develop novel probiotic formulations tailored to specific market segments [3].

The safety and efficacy of probiotics in combating foodborne pathogens have been highlighted in various studies, emphasizing their potential as natural and sustainable alternatives to antibiotics [4]. Probiotics have been investigated for their ability to enhance the shelf life, stability, and microbial safety of food products, thereby offering a viable solution to improving food preservation and quality. Moreover, the selection of probiotic strains based on their viability during processing and storage, as well as their ability to survive the gastrointestinal tract and exert health benefits, is crucial for their successful application in food systems [5].

This review aims to provide a comprehensive overview of the development of probiotics against food-borne bacterial pathogens, highlighting their mechanisms of action, efficacy, challenges, and future prospects.

2. MECHANISMS OF ACTION

Probiotics exert their antimicrobial effects through various mechanisms, including competitive exclusion, production of antimicrobial substances such as bacteriocins and organic acids, and modulation of the host immune system. Competitive exclusion involves the colonization of the gut mucosa by probiotic bacteria, preventing the attachment and subsequent colonization of pathogenic bacteria. Additionally, probiotics can produce antimicrobial substances that inhibit the growth of food-borne pathogens, thereby reducing their pathogenicity. Furthermore, probiotics interact with the host immune system, enhancing mucosal barrier function and promoting the production of antimicrobial peptides and immunoglobulins.

The development of probiotics against foodborne bacterial pathogens and understanding their mechanisms of action is a crucial area of research aimed at improving food safety and public health. Probiotics, such as *Lactobacillus* and *Bifidobacterium* species, have been extensively studied for their ability to combat foodborne pathogens like *Escherichia coli* and methicillin-resistant *Staphylococcus aureus* (MRSA) [1]. These probiotics have demonstrated significant antimicrobial activity against various pathogens, offering a promising approach to reducing the prevalence of harmful bacteria in food products and preventing foodborne illnesses. Moreover, probiotics have been found to modulate the immune system, maintain the mucosal barrier, and prevent conditions such as food allergies and diarrhoea.

Research has also explored the incorporation of probiotics into food packaging materials to control foodborne microorganisms and enhance food safety [2]. By encapsulating probiotics in biopolymer matrices, active food packaging materials have been developed as an innovative approach to extending the shelf life of food

products and providing health benefits. Additionally, the utilization of probiotics in both dairy and non-dairy products has been investigated to cater to consumer preferences and develop specialized probiotic formulations for different market segments [3].

The safety and efficacy of probiotics in combating foodborne pathogens have been underscored in various studies, highlighting their potential as natural and sustainable alternatives to traditional antibiotics [4]. Probiotics have been shown to enhance the shelf life, stability, and microbial safety of food products, offering a viable solution for improving food preservation and quality. Furthermore, the selection of probiotic strains based on their viability during processing and storage, as well as their ability to survive the gastrointestinal tract and confer health benefits, is critical for their successful application in food systems [5].

In recent years, there has been a growing body of literature focusing on the mechanisms of action of probiotics against foodborne bacterial pathogens. Studies have elucidated various mechanisms through which probiotics exert their beneficial effects, including the inhibition of pathogen growth, direct bactericidal effects, interference with virulence gene expression, competitive exclusion of pathogens, and stimulation of host immune responses [6]. These mechanisms collectively contribute to the antimicrobial activity of probiotics and their ability to combat foodborne pathogens effectively.

Moreover, research has highlighted the potential of probiotics to target specific multidrug-resistant bacteria, such as methicillin-resistant *Staphylococcus aureus* (MRSA) and vancomycin-resistant *Enterococcus* (VRE), underscoring the importance of probiotics as a strategy to address antibiotic resistance. Engineered probiotic strains have been proposed as a means to selectively target foodborne pathogens and inhibit their proliferation, offering a novel approach to food safety and public health [7].

Furthermore, the use of probiotics to modulate quorum sensing activity in foodborne pathogenic bacteria has emerged as a promising avenue for controlling bacterial virulence and communication, thereby enhancing food safety and gut health. Probiotic yeast metabolites have also been investigated for their cross-kingdom inhibition of bacterial virulence and communication, shedding light on the intricate interactions between probiotics and pathogenic bacteria [8].

3. EFFICACY AGAINST FOOD-BORNE PATHOGENS

Numerous studies have demonstrated the efficacy of probiotics against a range of food-borne pathogens. Probiotic strains such as *Lactobacillus* spp., *Bifidobacterium* spp., and *Enterococcus* spp. have shown promising results in reducing the colonization and shedding of pathogens such as *Salmonella*, *Escherichia coli*,

Listeria monocytogenes, and *Campylobacter jejuni*. These effects are attributed to the competitive exclusion of pathogens, production of antimicrobial compounds, and modulation of host immune responses. Moreover, probiotics have been investigated for their ability to mitigate the adverse effects of bacterial toxins and promote intestinal barrier integrity.

The development of probiotics as a strategy to combat food-borne bacterial pathogens has gained significant attention in recent years. Probiotics, defined as live microorganisms that confer health benefits when administered in adequate amounts, have shown promise in inhibiting the growth and activity of various food-borne pathogens [6]. Research has highlighted the efficacy of probiotics, particularly strains of *Lactobacillus* and *Bifidobacterium*, in combating bacterial pathogens such as *Escherichia coli*, *Salmonella*, *Listeria monocytogenes*, and *Staphylococcus aureus* [1, 9]. These probiotic strains have demonstrated antibacterial activities against a range of pathogens, showcasing their potential as biotherapeutic agents in addressing food-borne illnesses [1].

Studies have emphasized the importance of probiotics in preventing the colonization of pathogenic bacteria in the gastrointestinal tract by adhering to intestinal epithelial cells and stimulating host immunity [10]. Furthermore, probiotics have been shown to produce antimicrobial substances that inhibit the growth of food-borne pathogens, thereby enhancing food safety [11] [12]. The antagonistic effects of probiotics against a variety of food-borne pathogens, including *Escherichia coli*, *Salmonella*, *Listeria monocytogenes*, and *Bacillus cereus*, have been documented, underscoring their potential in controlling the proliferation of harmful bacteria [13].

Probiotics offer a multifaceted approach to combating food-borne pathogens, with mechanisms such as the production of organic acids, antimicrobial substances, and activation of the host's immune system contributing to their effectiveness [12]. Additionally, probiotics have been investigated for their ability to form biofilms that can inhibit the formation of biofilms by pathogens like *Listeria monocytogenes*, *Salmonella typhimurium*, and *Escherichia coli* O157:H7 [14]. This biofilm-forming capacity represents a novel strategy in the fight against food-borne pathogens, highlighting the versatility of probiotics in food safety applications.

The antimicrobial activities of probiotics and their metabolites have been extensively studied, showcasing their potential in inhibiting the growth of pathogenic microorganisms [15]. Probiotics have been found to produce bacteriocins, antimicrobial peptides that can serve as natural preservatives in food products, reducing the need for chemical additives and enhancing the nutritional quality of foods [16]. Moreover, the probiotic potential of lactic acid bacteria isolated from various sources has been explored, revealing their ability to inhibit

the growth of food-borne pathogens and improve intestinal microbial balance [17].

In the context of food preservation, probiotics have been investigated for their role in extending the shelf life of perishable products. Studies have demonstrated that probiotics can enhance the antioxidant activity of food matrices and exert antimicrobial effects against spoilage and pathogenic bacteria, thereby prolonging the freshness and safety of food products [18]. By incorporating probiotics into food packaging materials, researchers have explored innovative ways to enhance food quality and safety through natural means.

Overall, the development of probiotics against food-borne bacterial pathogens represents a promising avenue in ensuring food safety and public health. Through their diverse antimicrobial mechanisms, probiotics offer a natural and effective strategy to combat the proliferation of harmful bacteria in food products. Continued research into the efficacy, safety, and application of probiotics in food systems is essential for harnessing their full potential in mitigating the risks associated with food-borne illnesses.

4. CHALLENGES AND CONSIDERATIONS

Despite their potential, the development and application of probiotics in food safety face several challenges. Strain selection is crucial, as not all probiotic strains exhibit antimicrobial activity against food-borne pathogens. Furthermore, ensuring the viability and stability of probiotics during food processing and storage remains a challenge. Regulatory considerations, including safety and labelling requirements, also impact the commercialization of probiotic products. Additionally, the complexity of the gut microbiota and individual variability in probiotic response necessitate further research to optimize probiotic interventions.

The development of probiotics as a strategy against food-borne bacterial pathogens presents a promising avenue in enhancing food safety and human health. Probiotics, defined as live microorganisms that confer health benefits when administered in adequate amounts, have gained significant attention due to their potential to combat food-borne pathogens [19]. These beneficial bacteria, commonly including *Lactobacillus* and *Bifidobacterium* species, have been extensively studied for their ability to inhibit the growth of various food-borne pathogens such as *Escherichia coli*, *Salmonella enteritidis*, *Listeria monocytogenes*, and *Staphylococcus aureus* [13] [20]. Research has shown that probiotics can exert inhibitory effects on these pathogens through mechanisms such as antimicrobial compound production and competitive exclusion [20].

Incorporating probiotics into food products, such as whey cheese, has been explored as a means to reduce the risk of microbial contamination and enhance food safety [21]. Studies have demonstrated that probiotic strains like

Bifidobacterium animalis and *Lactobacillus casei*, when added to food matrices, exhibit antimicrobial activity against a spectrum of food-borne pathogens and spoilage microorganisms [21]. This application of probiotics in food production not only extends the shelf life of products but also contributes to the overall safety of the food supply chain.

Furthermore, the antimicrobial properties of probiotics have been linked to their ability to produce organic acids during fermentative metabolism, leading to antagonistic activities against bacterial pathogens by inducing intracellular acidification and membrane permeabilization [6]. These mechanisms showcase the potential of probiotics as natural agents for combating food-borne pathogens and highlight their role in promoting gastrointestinal health and preventing infections.

In addition to their direct antimicrobial effects, probiotics have been investigated for their anti-quorum sensing activity, which involves disrupting bacterial communication systems and inhibiting the virulence of pathogens. By interfering with quorum sensing, probiotics can impede the coordinated behaviour of pathogenic bacteria, thereby reducing their pathogenicity and enhancing food and gut health.

Moreover, the use of probiotics in combination with other strategies, such as paraprobiotics (sterilized probiotics) and postbiotics, has been proposed as a comprehensive approach to combat food-borne pathogens. These multifaceted interventions leverage the diverse mechanisms of probiotics, paraprobiotics, and postbiotics to target pathogens at different stages, from inhibition of growth to modulation of virulence factors.

The potential of probiotics to serve as biotherapeutic agents against gastrointestinal infections has been recognized, with studies highlighting their role in modulating the immune response and promoting gut health [9]. Probiotics have been shown to interact with the innate and acquired immune systems, offering a holistic approach to combating viral infections and other microbial threats [22].

Furthermore, the safety and efficacy of probiotics isolated from various sources, including fermented food products and traditional medicine, have been extensively evaluated [17, 23]. These studies emphasize the importance of selecting probiotic strains with proven antimicrobial activity against a broad spectrum of food-borne pathogens to ensure their beneficial impact on human health.

5. FUTURE DIRECTIONS

Future research efforts should focus on elucidating the mechanisms of action of probiotics against food-borne pathogens and identifying novel probiotic strains with enhanced antimicrobial properties. Strategies to improve the viability and stability of probiotics in food products are essential for their widespread application. Furthermore,

studies investigating the interactions between probiotics and the host immune system can provide valuable insights into their therapeutic potential. Integration of probiotics into food production and processing practices holds promise for enhancing food safety and reducing the risk of food-borne illnesses.

The development of probiotics against food-borne bacterial pathogens represents a crucial area of research with significant implications for food safety and human health. Probiotics, defined as live microorganisms that confer health benefits when administered in adequate amounts, have shown promise in combating food-borne pathogens. These beneficial bacteria exhibit antimicrobial activity against various foodborne pathogens, thereby reducing the risk of foodborne illnesses. Notably, *Lactobacillus* and *Bifidobacterium* species are commonly used probiotics that have demonstrated activity against diarrheagenic *Escherichia coli* (DEC) and other foodborne pathogens [1]. Additionally, probiotics like *Lactocaseibacillus* spp. have been found to possess antimicrobial and anti-biofilm activities against methicillin-resistant *Staphylococcus aureus* (MRSA).

Probiotics offer a multifaceted approach in combating food-borne pathogens, not only through direct antimicrobial effects but also by producing antimicrobial peptides that inhibit the growth of pathogens like *Escherichia coli* O157 H7, *S. aureus*, MRSA, and *Salmonella* Typhi. Furthermore, probiotics have been explored for their potential in developing active edible films and coatings, which could serve as innovative strategies for enhancing food safety by controlling foodborne microorganisms (Espitia et al., 2016). Encapsulation of probiotics into biopolymer matrices has led to the creation of active food packaging materials that can extend food shelf life and improve overall food safety.

The use of probiotics in food systems extends beyond direct consumption to applications in food packaging, where probiotics have been incorporated into biopolymeric materials to control foodborne pathogens and enhance food safety. Moreover, the impact of probiotics on the gut microbiota is a critical aspect to consider, as these beneficial microorganisms can modulate the intestinal microbial ecosystem, thereby influencing overall health [24]. Understanding the safety of probiotics is paramount, and research has focused on providing an evidence-based perspective on probiotic safety, particularly in probiotic fruit beverages and nutraceuticals [25].

Probiotics have also been investigated for their potential role in bio-preservation of plant-based foods and beverages, highlighting the importance of probiotics in enhancing food safety and extending the shelf life of perishable products. The future of probiotics lies in their continued development to meet consumer demands, including the creation of novel probiotic products tailored

to specific market segments such as infants, children, the elderly, and chronically ill patients. Additionally, advancements in probiotic research have paved the way for the development of next-generation probiotics with enhanced biotherapeutic properties.

6. CONCLUSION

Probiotics offer a promising approach for combating food-borne bacterial pathogens, providing an alternative to traditional antimicrobial therapies. Their ability to exert antimicrobial effects, modulate host immune responses, and promote intestinal barrier integrity makes them attractive candidates for food safety interventions. However, challenges such as strain selection, viability, stability, and regulatory considerations must be addressed to realize the full potential of probiotics in mitigating the risks associated with food-borne illnesses. Continued research efforts and collaborations between academia, industry, and regulatory agencies are essential for advancing the development and application of probiotics in food safety and public health.

The development of probiotics as a strategy against food-borne bacterial pathogens has gained significant attention in recent years. Probiotics, defined as live microorganisms that confer health benefits when administered in adequate amounts, have shown promise in combating various food-borne pathogens. *Lactobacillus* and *Bifidobacterium* species are commonly used probiotics that have demonstrated activity against diarrheagenic *Escherichia coli* (DEC) and other food-borne pathogens [1]. These probiotics produce antimicrobial substances such as bacteriocins, hydrogen peroxide, organic acids, and short-chain fatty acids, which inhibit the growth of bacterial and fungal pathogens by competing for resources and inhibiting pathogen adherence [26].

Studies have highlighted the ability of probiotics to restore balance to gut microbiota by outcompeting pathogenic bacteria and secreting antimicrobials to eliminate bacterial pathogens [27]. Additionally, engineered probiotic strains have been proposed as a potential intervention against food-borne pathogens, with the ability to selectively target and inactivate these pathogens, even in complex food matrices [7]. Probiotic lactobacilli, in particular, have been recognized for their antagonistic activity against pathogenic bacteria through the production of lactic acid, organic acids, hydrogen peroxide, and bacteriocins, creating an environment unfavourable for the growth of pathogenic bacteria.

The effectiveness of probiotics in combating food-borne pathogens is influenced by various factors, including the diversity of bacterial strains in probiotic products. Research has shown that probiotic products with a greater diversity of bacterial strains exhibit better antimicrobial activity against common gastrointestinal pathogens such as

Escherichia coli, *Shigella*, *Salmonella* spp., and *Clostridioides difficile*. Moreover, the use of probiotics as a biocontrol strategy to suppress pathogens in the gut has been proposed as a potential approach to managing bacterial infections.

Probiotics have also been explored for their potential in preventing dental caries by controlling the bacterial agents responsible for tooth decay [28]. Furthermore, the incorporation of probiotics into food packaging materials has been investigated as a means to control food-borne pathogens, enhance food safety, and provide health benefits. Probiotics have been utilized in various food formulations, including beverages, to leverage their functional properties and potential health benefits [29].

REFERENCES

1. Rosa ACP. Diarrheagenic *Escherichia Coli* and Probiotic Activity Against Foodborne Pathogens: A Brief Review. *Gastroenterology & Hepatology Open Access*. 2017;7(5); doi: 10.15406/ghoa.2017.07.00248.
2. Espitia PJP, Batista RA, Henriette Monteiro Cordeiro de A, Otoni CG. Probiotics and Their Potential Applications in Active Edible Films and Coatings. *Food Research International*. 2016; 90:42-52. doi: 10.1016/j.foodres.2016.10.026.
3. Ranadheera CS, Vidanarachchi JK, Rocha RS, Cruz AG, Ajlouni S. Probiotic Delivery Through Fermentation: Dairy vs. Non-Dairy Beverages. *Fermentation*. 2017 ;3(4):67; doi: 10.3390/fermentation3040067.
4. Reuben RC, Elghandour MM, Alqaisi O, Cone JW, Márquez O, Salem AZ. Influence of Microbial Probiotics on Ruminant Health and Nutrition: Sources, Mode of Action and Implications. *Journal of the Science of Food and Agriculture*. 2021;102(4):1319-40; doi: 10.1002/jsfa.11643.
5. Téptou A, Papadaki A, Lappa IK, Kachrimanidou V, Bosnea L, Kopsahelis N. Probiotics in Food Systems: Significance and Emerging Strategies Towards Improved Viability and Delivery of Enhanced Beneficial Value. *Nutrients*. 2019;11(7):1591; doi: 10.3390/nu11071591.
6. Moal VL, Servin AL. Anti-Infective Activities of *Lactobacillus* Strains in the Human Intestinal Microbiota: From Probiotics to Gastrointestinal Anti-Infectious Biotherapeutic Agents. *Clinical Microbiology Reviews*. 2014;27(2):167-99. doi: 10.1128/cmr.00080-13.
7. Yan X, Gurtler JB, Fratamico PM, Hu J, Juneja VK. Phylogenetic Identification of Bacterial MazF Toxin Protein Motifs Among Probiotic Strains and Foodborne Pathogens and Potential Implications of Engineered Probiotic Intervention in Food. *Cell & Bioscience*. 2012;2(1):39. doi: 10.1186/2045-3701-2-39.
8. Malka O, Kalson D, Yaniv K, Shafir R, Rajendran M, Ben-David O, et al. Cross-kingdom inhibition of bacterial virulence and communication by probiotic yeast metabolites. *Microbiome*. 2021;9(1):70. doi: 10.1186/s40168-021-01027-8.
9. Li P, Gu Q, Zhou Q. Complete Genome Sequence of *Lactobacillus Plantarum* LZ206, a Potential Probiotic Strain with Antimicrobial Activity Against Food-Borne Pathogenic Microorganisms. *Journal of Biotechnology*. 2016; 238:52-5; doi: 10.1016/j.jbiotec.2016.09.012.

10. Bian X, Evvie SE, Zafarullah M, Luo G, Liang H-z, Wang N, et al. In Vitro Assessment of the Antimicrobial Potentials of Lactobacillus Helveticus Strains Isolated from Traditional Cheese in Sinkiang China Against Food-Borne Pathogens. Food & Function. 2016;7(2):789-97. doi: 10.1039/c5fo01041a.
11. Adeniran AH, Ukeyima MT. Microbiological Assessment of <i>probioticated</i> Ginger-based Beverages. Nutrition & Food Science. 2010;40(2):209-20. doi: 10.1108/00346651011029246.
12. Karimi S, Rashidian E, Birjandi M, Mahmoodnia L. Antagonistic Effect of Isolated Probiotic Bacteria from Natural Sources Against Intestinal Escherichia Coli Pathotypes. Electronic Physician. 2018;10(3):6534-9; doi: 10.19082/6534.
13. Yazid AM, Salina A, Mustafa S, Osman H, Normah J. Inhibitory Effects of Probiotic Bacteria Against Selected Food-Borne Pathogens. Pakistan Journal of Biological Sciences. 1999;2(3):660-3; doi: 10.3923/pjbs.1999.660.663.
14. Gómez NC, Ramiro JMP, Quecán BXV, Bernadette Dora Gombossy de Melo F. Use of Potential Probiotic Lactic Acid Bacteria (LAB) Biofilms for the Control of Listeria Monocytogenes, Salmonella Typhimurium, and Escherichia Coli O157:H7 Biofilms Formation. Frontiers in Microbiology. 2016;7; doi: 10.3389/fmicb.2016.00863.
15. Sayeed MA, Hossen A, Saha RK, Jakaria M. Antimicrobial Activities of Isolated Probiotics and Their Metabolites Against Some Pathogenic Microorganisms. Iiuc Studies. 2018;14(1):21-8; doi: 10.3329/iiucs.v14i1.37652.
16. Das D, Goyal A. Characterization of a Noncytotoxic Bacteriocin from Probiotic Lactobacillus Plantarum DM5 With Potential as a Food Preservative. Food & Function. 2014;5(10):2453-62; doi: 10.1039/c4fo00481g.
17. Gupta AK, Sharma N. Probiotic Potential of Lactic Acid Bacteria Ch-2 Isolated from Chuli Characterization of Potential Probiotic Lactic Acid Bacteria- Pediococcus Acidilactici Ch-2 Isolated from Chuli- A Traditional Apricot Product of Himalayan Region for the Production of Novel Bioactive Compounds with Special Therapeutic Properties. Journal of Food Microbiology Safety & Hygiene. 2017;02(01); doi: 10.4172/2476-2059.1000119.
18. Yasin R, Samiullah K, Fazal RM, Hussain S, Mahboob S, Al-Ghanim KA, et al. Combined Effect of Probiotics on Prolonging the Shelf Life of GIFT Tilapia Fillets. Aquaculture Research. 2020;51(12):5151-62; doi: 10.1111/are.14853.
19. Goktepe I, Juneja VK, Ahmedna M. Probiotics in food safety and human health. 2005.
20. Sipahi N. Investigation of Antagonistic Effect of Probiotic Food Supplement Against Different Strains of Bacteria. International Journal of Traditional and Complementary Medicine Research. 2021;02(02):95-100. doi: 10.53811/ijtcmr.893132.
21. Madureira AR, Pintado ME, Gomes AMP, Malcata FX. Incorporation of Probiotic Bacteria in Whey Cheese: Decreasing the Risk of Microbial Contamination. Journal of Food Protection. 2011;74(7):1194-9. doi: <https://doi.org/10.4315/0362-028X.JFP-10-217>.
22. Kanauchi O, Andoh A, AbuBakar S, Yamamoto N. Probiotics and Paraprobiotics in Viral Infection: Clinical Application and Effects on the Innate and Acquired Immune Systems. Curr Pharm Des. 2018;24(6):710-7. doi: 10.2174/1381612824666180116163411.
23. Shivangi S, Devi PB, Ragul K, Shetty PH. Probiotic Potential of Bacillus Strains Isolated from an Acidic Fermented Food Idli. Probiotics Antimicrob Proteins. 2020;12(4):1502-13; doi: 10.1007/s12602-020-09650-x.
24. Gargari G, Taverniti V, Koirala R, Gardana C, Guglielmetti S. Impact of a Multistrain Probiotic Formulation with High Bifidobacterial Content on the Fecal Bacterial Community and Short-Chain Fatty Acid Levels of Healthy Adults. Microorganisms. 2020;8(4):492. doi: 10.3390/microorganisms8040492.
25. Žuntar I, Petrić Z, Kovačević DB. Safety of Probiotics: Functional Fruit Beverages and Nutraceuticals. Foods. 2020;9(7):947. doi: 10.3390/foods9070947.
26. Li P, Gu Q. Antimicrobial Effects of Probiotics and Novel Probiotic-Based Approaches for Infectious Diseases. 2018; doi: 10.5772/intechopen.72804.
27. Hillman T. Quantifying the Antimicrobial Activity of CRISPR-Cas9-accA Modified ΔB. Subtilis mutants Against V. Harveyi and E. Coli. 2021. doi: 10.1101/2021.08.10.455802.
28. Shakib P, Rouhi S, Zolfaghary MR. The Role of Probiotics in Preventing Dental Caries. Plant Biotechnology Persa. 2020;2(1):55-8; doi: 10.29252/pbp.2.1.55.
29. Sharma S, Singh A, Sharma S, Kant A, Sevda S, Taherzadeh MJ, et al. Functional Foods as a Formulation Ingredients in Beverages: Technological Advancements and Constraints. Bioengineered. 2021;12(2):11055-75. doi: 10.1080/21655979.2021.2005992.