

## Simple and Inexpensive Methods Development for the estimation of Tamsulosin Hydrochloride as a single component from Its Solid Dosage Forms by Visible Spectrophotometry

K. Raghubabu<sup>1</sup>, L. Shanti swarup<sup>1</sup>, B. Kalyanaramu<sup>\*2</sup>, M. N. Rao<sup>3</sup>, C. Ramdas<sup>3</sup>

<sup>1</sup>Department of Engineering Chemistry, AU College of Engineering (A), Andhra University, Visakhapatnam-530003, AP (India)

<sup>2</sup>Department of Chemistry, M. R. College for Men (A), Vizianagaram-535002, AP (India).

<sup>3</sup>M/s Tychy Industries, Hyderabad, AP (India)

\*Corresponding Author Email: [Kalyanaramu23566@gmail.com](mailto:Kalyanaramu23566@gmail.com)

**BIOLOGICAL SCIENCES**

**Research Article**

RECEIVED ON 02-12-2011

ACCEPTED ON 12-1-2012

### ABSTRACT

Tamsulosin hydrochloride (TAM) is an uroselective  $\alpha_{1A}$ -adrenergic receptor antagonist which is used in benign prostatic hyperplasia (BPH). The drug is official in European Pharmacopoeia. Two direct, simple and sensitive visible spectrophotometric methods are described for the assay of tamsulosin hydrochloride in pure and solid dosage forms. The first method  $M_1$  is based on the formation of yellowish brown colored species by the drug with Folin reagent and exhibits absorption maxima at 440 nm. Second method ( $M_2$ ) is based on the formation of purple red colored species with sodium nitroprusside-acetaldehyde reagent exhibiting maximum absorption at 560 nm. Regression analysis of Beer-Lambert plots showed good correlation in the concentration ranges (16-48)  $\mu\text{g/ml}$  for method  $M_1$ , and (8.0-24)  $\mu\text{g/ml}$  for method  $M_2$  respectively. The proposed methods are applied to commercial available tablets and the results are statistically compared with those obtained by the UV reference method and validated by recovery studies. The results are found satisfactory and reproducible. These methods are applied successfully for the estimation of the tamsulosin hydrochloride in the presence of other ingredients that are usually present in dosage forms. These methods offer the advantages of rapidity, simplicity and sensitivity and normal cost and can be easily applied to resource-poor settings without the need for expensive instrumentation and reagents.

**KEYWORDS:** Assay, BPH, Folin reagent, Sodium nitroprusside, Acetaldehyde, Nucleophilic substitution, Statistical analysis.

### INTRODUCTION

Tamsulosin hydrochloride (TAM) (Fig.1), 5-[(2R)-2-[2-(2-ethoxyphenoxy) ethylamino] propyl]-2-methoxybenzenesulfonamide hydrochloride is a new type of highly selective  $\alpha_{1A}$  adrenergic receptor antagonist for treatment of benign prostatic hyperplasia (BPH)<sup>1</sup>. The drug exists in two enantiomeric forms, but only R-isomer is the pharmaceutically active component. It works by blocking  $\alpha$ -receptors that are found in the muscle of the prostate gland, which causes the muscle in

the prostate to relax. This allows urine to flow freely past the prostate and relieves the urinary symptoms. The drug is extensively metabolized by cytochrome P450 enzymes in the liver and the most frequently prescribed medication for the treatment of lower urinary tract symptoms associated with BPH. Compared to other  $\alpha$ -antagonists, TAM has greater specificity for  $\alpha_1$ -receptors in the human prostate and does not affect receptors on blood vessels. The drug is official in European Pharmacopoeia<sup>2</sup>.




Fig.1: Chemical structure of TAM

Some analytical methods which include HPLC<sup>3-11</sup>, LC-MS-MS<sup>12-18</sup>, HPTLC<sup>19-21</sup> Radioreceptor assay<sup>22-23</sup>, Non aqueous potentiometric<sup>24</sup>, voltametry<sup>25</sup>, Capillary electrophoresis<sup>26</sup>, spectrofluorimetric<sup>27</sup> UV<sup>28-29</sup> and visible spectrophotometric<sup>30-31</sup> have been reported in the literature for Determination of TAM in biological fluids (more) and pharmaceutical preparations(less). The main purpose of the present study was to establish relatively simple, sensitive, validated and inexpensive extraction free Visible spectrophotometric methods for the determination of TAM in pure form and in pharmaceutical preparations, since most of the previous methods involve critical reaction conditions or tedious sample preparations and less specificity. So the authors have made some attempts in this direction and succeeded in developing these methods based on the reaction between the drug and folin reagent<sup>32</sup> (M<sub>1</sub>) or sodium nitro prusside-acetaldehyde reagent<sup>33</sup> (M<sub>2</sub>) under specified experimental conditions. The proposed methods for TAM determination have many advantages over other analytical methods due to its rapidity, normal cost and environmental safety. Unlike HPLC, HPTLC procedures, the instrument is simple and is not costly. Economically, all the analytical reagents are inexpensive and available in any analytical laboratory. These methods can be extended for the routine quality control analysis of pharmaceutical products containing TAM.

## MATERIALS & METHODS (EXPERIMENTAL)

**Apparatus and chemicals:** A Shimadzu UV-Visible spectrophotometer 1601 with 10mm matched quartz cells was used for all spectral measurements. A Systronics digital pH meter mode-361 was used for pH measurements. All the chemicals used were of analytical grade. Pure TAM drug was obtained as a gift sample from M/s Tychy Industries, Hyderabad (AP). Tablets were purchased from local market.

Folin reagent (NQS) solution (Loba, 0.5%, 1.92x10<sup>-2</sup>M prepared by dissolving 500mg of NQS in 100 ml of distilled water), aqueous solutions of sodium nitro prusside (SNP, E. Merck, 1.0%, 3.35x10<sup>-2</sup>M), acetaldehyde (10%), phosphate buffer of pH 8.0

(prepared by mixing 30 ml of 0.067M potassium hydrogen phosphate and 970 ml of 0.067M disodium hydrogen phosphate and pH adjusted to 8.0) were prepared.

**Preparation of Standard stock solution:** The standard stock solution (1mg/ml) of TAM was prepared by dissolving 100mg of TAM in 10 ml 0.1M sodium hydroxide and the volume was brought to 100 ml with distilled water. The working standard solutions of TAM were obtained by appropriately diluting the standard stock solution with the same solvent (M<sub>1</sub>- 400 µg/ml & M<sub>2</sub>-200 µg/ml). The prepared stock solution was stored at 4<sup>o</sup> C protected from light. From this stock solution, a series of standards were freshly prepared during the analysis day.

**Preparation of Sample solution:** About 20 tablets or capsules were weighed to get the average tablet or capsule weight and pulverized. The powder equivalent to 100mg of TAM was weighed, dispersed in 25ml of Isopropyl alcohol, sonicated for 15 minutes and filtered through Whatman filter paper No 41. The filtrate was evaporated to dryness and the residue was dissolved as under standard solution preparation.

### Determination of wavelength maximum (λ<sub>max</sub>):

**Method M<sub>1</sub>:** The 3.0 ml of working standard solution of TAM (400µg/ml) was taken in 25ml standard flask. To this, 1.0ml of folin reagent (1.092x10<sup>-2</sup>M), 5.0 ml of buffer pH 8.0 and 1.5ml of distilled water were added and kept aside for 15 min for complete color development. Then the volume was made up to 25 ml using distilled water and sonicated for 1 min. to get a concentration of 48µg/ml. In order to investigate the wavelength maximum, the above standard stock solution was scanned in the range of 360-560nm by UV-Visible spectrophotometer. From the spectra (**Fig.2**), it was concluded that 440nm is the most appropriate wavelength for analyzing TAM with suitable sensitivity.

**Method M<sub>2</sub>:** The 3.0 ml of working standard solution of TAM (200µg/ml) was taken in 25 ml calibrated tubes containing 15ml of buffer pH 8.0. To this, 1.0 ml each of SNP solution and acetaldehyde were added successively and shaken

for 2 minutes and kept aside for 15 minutes at room temperature and made up to the mark with distilled water and sonicated for 1 min, to get a concentration of 24µg/ml. In order to investigate the wavelength maximum, the colored solution was scanned in the range of 400-700nm by UV-Visible spectrophotometer. From the UV spectra (Fig.4), it was concluded that 560nm is the most appropriate wavelength for analyzing TAM with suitable sensitivity.

**Preparation of calibration curve:**

Aliquots of the standard TAM solution [1.0-3.0ml, 400µg/ml (M<sub>1</sub>)] were placed in a series of 25ml standard flask. Then 1.0ml of folin reagent (1.092x10<sup>-2</sup>M), 5.0 ml of buffer pH 8.0 and 1.5ml of distilled water were added and kept aside for 15 min for complete color development. Then the volume was made up to 25 ml using distilled water and sonicated for 1 min. The absorbance was measured at 440 nm (M<sub>1</sub>) against a reagent blank

within the stability period (5minutes to 30min). The calibration graph was constructed by plotting the drug concentration versus absorbance. The amount of drug was computed from its calibration graph (Fig. 3).

Aliquots of working standard TAM drug solution (200µg/ml) such as 1.0, 1.5, 2.0, 2.5, and 3.0 ml were taken separately in a series of 25ml calibrated tubes containing 15ml of buffer pH 8.0. Then 1.0ml each of SNP solution and acetaldehyde were added successively and shaken for 2 minutes and kept aside for 5 minutes at room temperature and made up to the mark with distilled water and sonicated for 1 min. The absorbance was measured at 560 nm (M<sub>2</sub>) against a reagent blank within the stability period (5minutes to 30min).The calibration graph was constructed by plotting the drug concentration versus absorbance. The amount of drug was computed from its calibration graph (Fig. 5).




Fig.2: Absorption spectra of TAM-NQS




Fig.3: Beer's Law plot of TAM-NQS




Fig.4: Absorption spectra of TAM-SNP-ACD




Fig.5: Beer's Law plot of TAM-SNP-ACD

## RESULTS AND DISCUSSION

Optimum operating conditions used in the procedure were established by adopting variation of one variable at a time (OVAT) method. The effect of various parameters such as time, volume and strength of reagents, the order of addition of reagents, pH buffer solutions and solvent for final dilution of the colored species were studied. Distilled water was found to be best solvent for final dilution. Other water miscible solvents like methanol, ethanol, propan-2-ol and acetonitrile have no additional advantage in increasing the intensity of the color in both methods. The optical characteristics such as Beer's law limit, Sandell's sensitivity, molar absorptivity, percent relative standard deviation, (calculated from the six measurements containing 3/4<sup>th</sup> of the amount of the upper Beer's law limits), Regression characteristics like standard deviation of slope ( $S_b$ ), standard deviation of intercept ( $S_a$ ), standard error of estimation ( $S_e$ ) and % range of error (0.05 and 0.01 confidence limits) were calculated and the results are summarized in **Table-1**.

Commercial formulations containing TAM were successfully analyzed by the proposed methods. The values obtained by the proposed and reference methods for formulations were compared statistically by the t-and F-test and found not to differ significantly. As an additional demonstration of accuracy, recovery experiments were performed by adding a fixed amount of the drug to the pre analyzed formulations at three different concentration levels. These results are summarized in **Table-2**.

**TABLE 1: OPTICAL CHARACTERISTICS, PRECISION AND ACCURACY OF PROPOSED METHODS**

Parameter	Method M <sub>1</sub>	Method M <sub>2</sub>
$\lambda_{\max}$	440 nm	560 nm
Beer's law limit( $\mu\text{g/ml}$ )	16-48	8-24
Sandell's sensitivity ( $\mu\text{g/cm}^2/0.001$ abs. unit)	0.004238411	<b>0.002191781</b>
Molarabsorptivity (Litre/mole/cm)	104987.4688	<b>203022.125</b>
Correlation coefficient Regression equation (Y)*	0.998	<b>0.999</b>
Intercept (a)	-0.133	<b>-0.112</b>
Slope(b)	0.013	<b>0.025</b>
%RSD	1.114	<b>0.927</b>
% Range of errors(95% Confidence limits)		
<b>0.05 significance level</b>	1.17	<b>0.973</b>
<b>0.01 significance level</b>	<b>1.83</b>	<b>1.53</b>

\*Y = a + b x, where Y is the absorbance and x is the concentration of TAM in  $\mu\text{g/ml}$ .

**TABLE-2: ANALYSIS OF TAMSULOSIN HYDROCHLORIDE IN PHARMACEUTICAL FORMULATIONS BY PROPOSED AND REFERENCE METHODS.**

Method	*Formulations	Labeled Amount (mg)	Found by Proposed Methods			Found by Reference Method $\pm$ SD	#% Recovery by Proposed Method $\pm$ SD
			**Amount found $\pm$ SD	t	F		
A	Batch-1	0.2	0.199 $\pm$ 0.001	0.513	4.57	0.199 $\pm$ 0.0007	99.54 $\pm$ 0.73
	Batch-2	0.4	0.398 $\pm$ 0.002	0.38	3.72	0.397 $\pm$ 0.0044	99.41 $\pm$ 0.56
B	Batch-1	0.2	0.199 $\pm$ 0.0007	0.164	1.32	0.199 $\pm$ 0.0007	99.406 $\pm$ 0.389
	Batch-2	0.4	0.398 $\pm$ 0.002	0.95	3.87	0.397 $\pm$ 0.0044	99.59 $\pm$ 0.553

\* Different batches from two different companies \*\*Average  $\pm$  Standard deviation of six determinations, the t- and F-values refer to comparison of the proposed method with reference method (UV). Theoretical values at 95% confidence limits t = 2.57 and F = 5.05.

# Recovery of 10mg added to the pre-analyzed sample (average of three determinations). Reference method (reported UV method) using distilled water ( $\lambda_{\max}$ =279nm).

### Chemistry of colored species:

In the present investigation, the presence of aliphatic secondary amino group of TAM permits the development of visible spectrophotometric methods for its determination. In method M<sub>1</sub>, yellowish brown colored species (N-alkyl amino naphthaquinone) was formed by replacement of the sulphonate group of the naphthaquinone sulphonic acid by a secondary amino group of drug.

Cullies and Waddington<sup>34</sup> found that many secondary but not primary or tertiary amines react with sodium nitro prusside and acetaldehyde

under mild alkaline conditions. Wolfe and Swine hart<sup>35</sup> have reported the formation of  $[\text{Fe}(\text{CN})_5\text{H}_2\text{O}]^{3-}$  in aqueous solution of sodium nitro prusside. The proposed method M<sub>2</sub> exploits structural features aliphatic secondary amine of the TAM molecule. The nature of colored species formation with sodium nitro prusside-acetaldehyde reagent is initial N-alkyl vinyl amine formation with acetaldehyde then followed by formation of colored inner molecular complex with sodium nitro prusside. The formation of colored species with these reagents may be assigned through above analogy as shown in schemes (Fig.6).





## CONCLUSION

The reagents utilized in the proposed methods are cheap, readily available and the procedure does not involve any critical reaction conditions or tedious sample preparation. The proposed colorimetric methods possesses reasonable precision, accuracy, and are simple, sensitive and can be used as alternative methods to the reported ones for the routine determination of TAM depending on the need and situation.

## ACKNOWLEDGEMENTS

The authors are very much thankful to the m/s Tychy Industries, Hyderabad, Andhra Pradesh (India) for providing gift sample of the drug and also thanks to the University authorities for providing necessary facilities.

## REFERENCES

1. The Merck Index, Merck Research laboratories, Division of Merck Co Inc., Whitehouse station NJ, 2006, 1728.
2. European Pharmacopoeia, 5<sup>th</sup> ed. Suppl. 5.7, European Directorate for the quality of medicines; 2006, p.5121.
3. Kumari Richa, Dash PP, Lal VK, Mishra A, Murthy PN, RP-HPLC method for the estimation of tamsulosin hydrochloride in tablet dosage form. Indian Journal Pharmaceutical Science 2010; 72(6):785-787.
4. Sudha Thomas, Jitendra Dhomane, A validated RP-HPLC method for the determination of impurities in tamsulosin hydrochloride. Int J Chem Res 2011; 2(4): 29-33.
5. Chandorkar JG, Kotwal VB, Dhande NS, Gurav SG, Pande VV, Yadav PV, A sensitive HPLC method for simultaneous estimation of tamsulosin hydrochloride and its impurity. Pak J Pharm. Sci., 2008; 21: 307-310.
6. Zhang Z, Yang G, Liang G, Liu, H, Chen Y, Chiral separation of tamsulosin isomers by HPLC using cellulose Tris (3,5-dimethylphenylcarbamate) as a chiral stationary phase. J Pharm Biomed Anal 2004; 34(3): 689-93.
7. Qi M, Wang P, Cong R, Determination of the enantiomers of tamsulosin and its synthetic intermediates by chiral LC. Chromatographia 2004; 59(3-4):251-254.
8. Macek J, Klíma J, Ptacek P, Rapid determination of tamsulosin in human plasma by HPLC using extraction with butyl acetate. J Chromatogr B 2004; 809: 307-11.
9. Rao BM, Srinivasu MK, Sridhar G, Reddy BS, Vittal TV, Kumar RP et al. HPLC studies on the stress degradation behavior of tamsulosin hydrochloride and development of validated specific stability indicating method. Indian Drugs 2006; 43: 39-43.
10. Basaveswara Rao MV, Reddy BC, Rao TS, Rao VG, Drug release method by HPLC for tamsulosin hydrochloride 0.2% and Tolterodine tartrate 0.2% combination pellets. Int J Chem Eng Res 2009; 1: 155-159.
11. Bakshi M and Singh S, Development of validated stability indicating assay methods-critical review. Journal of Pharmaceutical and Biomedical Analysis 2002; 28(6):1011-1040.
12. Nageswra Rao R, Kumar Talluri MV, Narasa Raju A, Shinde DD, Ramanjaneyulu GS, Development of a validated RP-LC/ESI-MS-MS method for separation, identification and determination of related

substances of tamsulosin in bulk drug and formulations. J Pharm Biomed Anal 2008; 46(1): 94-103.

13. Ramakrishna NVS, Vishwottan KN, Manoj S, Koteswara M, Wishu S and Varma DP, Rapid simple and highly sensitive LC-ESI-MS/MS method for the quantification of tamsulosin in human plasma. Biomedical Chromatography 2005; 19(10): 709-719.
14. Keski-Rahkonen P, Parsinen O, Leppanen E, Mauriala T, Lehtonen M, Auriola S, Determination of tamsulosin in human aqueous humor and serum by liquid chromatography-electro spray ionization tandem mass spectrometry. J Pharm Biomed Anal 2007; 43(2): 606-12.
15. Ding Li, Li L, Tao P, Yang J, Zhang Z, Quantitation of tamsulosin in human plasma by liquid chromatography-electro spray ionization mass spectrometry. J. Chromatogr B 2002; 767(1):75-81.
16. Qi M, Wang P, Liu L, Determination of tamsulosin in dog plasma by liquid chromatography with atmospheric pressure chemical ionization tandem mass spectrometry J. Chromatogr B 2004; 805: 7-11.
17. Fan HR, Gu Y, Si DY, Liu CX, Determination of tamsulosin in dog plasma by a high sensitive liquid chromatography tandem mass spectrometric method. Yao Xue Xue Bao 2007; 42: 872-76.
18. Matsushima H, Takanuki KI, Kamimura H, Watanabe T and Higuchi S, Highly sensitive method for the determination of tamsulosin in human plasma dialysate, plasma and urine by HPLC-electro spray tandem mass spectrometry. Drug Metab. Dispos. 2004; 26: 240-245.
19. Bari SB, Bakshi AR, Jain PS and Surana SJ, Development and validation of stability indicating HPTLC determination of tamsulosin in bulk and pharmaceutical dosage form. Chromatography Research International 2011; Vol.2011:1-6.
20. Patel DB, Patel NJ, Validated stability indicating HPTLC method for the determination of tamsulosin hydrochloride in pharmaceutical dosage forms. Int J Chem Tech Res 2010; 2(1): 646-652.
21. Patel D and Patel N, Validated RP-HPLC and TLC methods for simultaneous estimation of tamsulosin and finasteride in combined dosage forms. Acta Pharmaceutica 2010; 60(2):197-205.
22. Taguchi K, Schafers RF, Michel MC, Radioreceptor assay analysis of tamsulosin and terazosin pharmacokinetics. Br. J Clin. Pharmacol. 1998; 45: 49-55.
23. Yamada S, Tanaka C, Suzuki M, Ohkura J, Kimura and Kawabe K, Determination of alpha- adrenoceptor antagonists in plasma by radioreceptor assay. Journal of Pharmaceutical and Biomedical Analysis 1996; 14(3):289-94.
24. Rao BM, Srinivasu MK, Thilakumar T, More S, Rajendra kumar P, A validated non-aqueous potentiometric titration method for the quantitative determination of tamsulosin in bulk drug. Indian Drugs 2005; 42:175-177.
25. Ozkan SA, Uslu B, Aboul-Enein HY, Voltammetric investigation of tamsulosin. Talanta 2003; 61: 147-56.
26. Maier V, Horakova J, Petr J, Tesarova C, Coufal P, Sevcik J, Chiral separation of tamsulosin by capillary electrophoresis. J Pharm Biomed Anal 2005; 39(3-4): 691-96.
27. Patel, Patel NU, and Chaudhari BG, Validated spectrofluorimetric method for the determination of amsulosin hydrochloride in tablet dosage form. Der Pharmacia Sinica; 2(3): 172-175.
28. Gadhawe Nilam A, Sawant SD, Ghante MR, Nikam AD, Spectrophotometric estimation of tamsulosin hydrochloride in tablet dosage form. International Journal of Pharmaceutical Research and Development 2011; 3(4): 87-92.
29. Nanda RK, Gaikwad J and Prakash A, Asian journal of Research Chem. 2009; 2(1): 63-65.
30. Shrivastava A, Saxena P, Gupta VB, Spectrophotometric estimation of tamsulosin hydrochloride by acid-dye method. Pharmeceutical methods 2011; 2(1): 53-59.
31. Susmitha K, Radha K and Venkateshwarlu G, Extractive spectrophotometric methods for determination of tamsulosin hydrochloride in pharmaceutical formulations using acidic tri phenyl methane dyes. Asian journal of Research in Chemistry 2011; 4(7): 1114-18.

32. Rosenblatt DH, Hlinka P, Epstein J. Colorimetric determinations of aromatic amines. *Anal Chem* 1955; 27: 1290.  
33. CSN Sarma, C.Kamala sastri and CSP Sastry, *Asian J. Chem.*, 2002; Vol.14 (2): 691-698.

34. CP.Cullis and B.J.Waddington, *Anal.Chem.Acta*, 1956; 15: 158.  
35. SK Wolfe and JH. Swinehart, *Inorg.Chem*, 1975; 14:1049



**\* Corresponding Author**

B.Kalyanaramu\*  
Department of Chemistry, M. R. College for Men (A),  
Vizianagaram-535002, AP (India).