

International Journal of Pharmacy and Biological Sciences ISSN: 2321-3272 (Print), ISSN: 2230-7605 (Online) IJPBSTM | Volume 8 | Issue 4 | OCT-DEC | 2018 | 962-968



Research Article | Biological Sciences | Open Access | MCI Approved ज्ञान-विज्ञान विमुक्तये

UGC Approved Journal

IN VITRO EVALUATION OF ANTI OXIDANT ACTIVITY, ESTIMATION OF TOTAL PHENOLIC AND FLAVONOID CONTENT OF DIFFERENT EXTRACTS OF *TECOMA STANS* (L.)

R. Prathibha^{1*}, A. Niranjan Kumar², KVNS. Srinivas², J. Kotesh Kumar², A. Srivani¹, G. Krishna Mohan¹

¹CPS, IST, Jawaharlal Nehru Technological University, Hyderabad, India. ²Natural Product Chemistry, CSIR-Central Institute of Medicinal and Aromatic Plants, Research Centre, Boduppal, Hyderabad-500092, India.

*Corresponding Author Email: ramisettiprathibha@gmail.com

ABSTRACT

Tecoma stans is a flowering perennial plant characterized by the presence of yellow colored trumpet-shaped flowers. The various parts of the plant possess several medicinal properties due to the presence of diversified group of biologically/pharmacologically active phytochemicals and are used therapeutically for the treatment of various diseases such as diabetes, cancer, yeast infections etc. The plant is reported to hold the medicinal value as a powerful diuretic, vermifuge, cardiotonic, anti-spasmodic agent apart from its use as an ornamental, fuel, timber, shade/shelter, and barrier. Based on the consideration of these facts, we engaged to study in-vitro antioxidant activity, determination of total phenolic and flavonoid content of different extracts of T.stans. The antioxidant activity was observed to be high in stem methanolic extract. The leaf methanolic extract tends to contain more phenolic content (60.24 mg GAE/ gr. Ext.). The total flavonoid content was observed to be high in leaf methanolic extract (127.18 mg CE/gr.Ext.) in terms of quercetin and catechin equivalents respectively.

KEY WORDS

Phytochemicals, Vermifuge, Cardiotonic, Anti-spasmodic, Catechin, Quercetin.

1. INTRODUCTION

The **Tecoma stans** plant is a speedy growing evergreen medicinal plant belonging to the family Bignoniaceae which grows to a height of about 20-30 feet and it is characterized by the presence of trumpet-shaped yellow colored flowers ^[1]. The plant is spread throughout the tropical and subtropical regions of India, South America, Florida, Bahamas, Trinidad, Africa, Asia and Australia ^[2]. The entire plant possesses medicinal value and the roots are used as diuretic, tonic, anti-syphilitic and vermifuge ^[3]. The decoction of flowers and bark are used for stomach pains. The plant is used in the treatment of diabetes ^[4]. The bark and flowers possess antimicrobial activity ^[5] and also used for the treatment

of various cancers ^[6]. The bark shows smooth muscle relaxant ^[7] and mild cardiotonic activity. The root of the plant is reported to be a powerful diuretic, vermifuge, and tonic ^[8]. A grinding of the root and lemon juice is reportedly used as a remedy for snake and rat bites. It is also reported to have cardioprotective action ^[9]. The whole plant has promising activities like Wound Healing Potential ^[10], Antispasmodic Effect ^[11], Antimicrobial Activity ^[12], Cytotoxic Activity ^[13,14], Antifungal Activity ^[15], Anti-Inflammatory Activity ^[16] and Anti-diabetic Activities due to the presence of various phytoconstituents like Sugars, Terpenoids, Phenolics and Alkaloids [17,18] The alkaloids Tecomine, Tecostanine, 5 Beta-Hydroxyskitanthine and Boschniakine isolated from T. stans plant exhibited good



anti-diabetic activity ^[19]. Apart from the medicinal usage, this plant was being widely used as an ornamental, fuel, timber, shade/shelter, barrier ^[20].

The plants rich in anti-oxidants are helpful for the prevention as well as the decrease in the oxidative stress and damage caused by the free radicals. The anti-oxidants inhibit the initiation and propagation of oxidative chain reactions. In view of the varied pharmacological and phytochemical reports of T. *stans,* the study on *in vitro* antioxidant activity and estimation of total phenolic and flavonoidal content of leaf and stem of *T. stans* was taken as part of our continuing phytochemistry work.

2. MATERIALS AND METHODS

2.1. Plant collection:

The leaf and stem parts of the plant *T. stans* were collected from CSIR-CIMAP, Research Centre, Hyderabad, Telangana, India. The plant was identified taxonomically and it was authenticated by Dr.A.Sabitha Rani, Assistant Professor, Department of Botany, Osmania University, Hyderabad, India. A voucher specimen (CIMAP-TS/18) was stored at CSIR-CIMAP Research Centre, Hyderabad.

2.2. Extraction:

The *T. stans* leaf and stem material (each 100 gm) were shade dried, powdered and successively extracted by the solvents with increasing order of polarity i.e Hexane, Ethyl acetate, Methanol by sonication equipment and also extracted with aq.methanol by Maceration method. Later, the extract solutions were filtered and distilled under reduced pressure using Rota vapor to obtain the crude extracts. These extracts were used for determination of Anti-oxidant activity and estimation of total phenolic & flavonoidal content.

Int J Pharm Biol Sci.

2.3. Determination of *In-Vitro* Anti-Oxidant Activity:

The ability of the different extracts of *T. stans* leaf and stem to scavenge DPPH radical was determined by using DPPH radical scavenging method ^[21]. DPPH is a darkcolored crystalline, nitrogen-centered substance composed of stable free radicals. The principle involves reduction of DPPH (purple color) in the presence of antioxidant in the sample which results in the formation of non-radical DPPH (yellow color). The color transformation of purple to yellow is measured spectrometrically at 517nm. The decrease in absorbance indicates greater inhibition of DPPH free radicals which is an indication of higher antioxidant content in the sample.

The sample extract stock solutions (1mg/mL) were diluted to the final concentrations of 5, 10, 15, 20, 25, 50 μ g/mL using methanol. To 5 mL of different extract solutions, 2 mL of 0.3 mM DPPH in methanol solution was added and allowed to react for 30 minutes at room temperature in a dark chamber. The reduction of DPPH radicals was measured by checking the absorbance in UV-VIS Spectrophotometer at 517nm.

Methanol (5mL) was used as the blank. DPPH solution (2mL) plus methanol (5mL) was used as negative control. The standard solutions of ascorbic acid 5, 10, 15, 20, 25, 50µg/mL were used as positive control.

The percentage inhibition of DPPH radicals by the antioxidant is calculated by using the formula-

Percentage of Inhibition (%) –	Blank absorbance - Sample absorbance	v	100
$\frac{1}{2} = \frac{1}{2} = \frac{1}$	Blank absorbance	л	100

2.4. Estimation of Total Phenolic content:

The total phenolic content of different extracts of *T. stans* leaf and stem was determined by Folin-Ciocalteus assay method ^[22]. The folin – cioaclteau reagent comprising of phosphotunstat–phosphomolybdenum complex (yellow) gets reduced by phenolate ion in the sample, changing its color to blue in the presence of a base. The color transformation is measured spectrometrically and the greater phenolic ion content in the sample is indicated by the increased absorbance. An aliquot (1 mL) of different extracts or standard solution of Gallic acid (50, 100, 150, 200 and 250 mg/lit) was added to 25 ml volumetric flask. To this flask, 9 mL

of distilled water was added. Further 1 mL of Folin-Ciocalteu phenol reagent was added to this flask and shaken for few minutes. To the mixture, 10 mL of 7 % sodium carbonate (Na₂CO₃) solution was added after 5 minutes. The solution mixture was diluted to 25 mL with distilled water. The solution was placed aside for incubation for about 90 minutes at room temperature. Similarly, a reagent blank solution was prepared by omitting the samples to be analyzed. The absorbance of the sample extract and standard solutions against the prepared reagent blank was observed at 750 nm with the UV-Visible spectrophotometer. The phenolic content expressed as mg Gallic Acid Equivalents (GAE)



per gram extract weight. The total phenolic content was calculated by using the formula-

Total Phenolic Content=	Concentration	x	Vol. of the Sample Weight of the Sample	
-------------------------	---------------	---	--	--

2.5. Estimation of Total Flavonoidal content:

The total flavonoidal content of different extracts of *T. stans* leaf and stem were determined by aluminium chloride colorimetric assay using Quercetin and Catechin as the standards. The principle is based on the formation of acid stable complexes by aluminium chloride with the keto group and hydroxyl group of flavonoids.

Colorimetric assay (Quercetin)

The total flavonoid content was measured by the aluminium chloride colorimetric assay using quercitin as standard ^[23]. An aliquot of 2.5 mL of extracts or standard solution of Quercitin (150, 200, 300, 400 and 500 mg/lit) was added to 25 mL volumetric flask. To this flask, 10 mL of distilled water was added. Further 0.75 mL of 5 % sodium nitrite (NaNO₂) was added. To this solution mixture, 0.75 mL of 10 % aluminium chloride (AlCl₃) was added after 5 minutes. At the 6th min, 2 mL of 1 M sodium hydroxide (NaOH) was added and the total volume was made up to 25 mL with distilled water.

Similarly, the reagent blank was prepared by omitting the samples to be analyzed. The solutions were mixed well, and the absorbance was measured against the prepared reagent blank at 510 nm in UV-VIS spectrophotometer. The flavonoid content was expressed as mg Quercetin equivalents/gr. weight of extract.

Colorimetric assay (Catechin):

The total flavonoid content was measured by the aluminium chloride colorimetric assay using catechin as

standard ^[24]. An aliquot of 2.5 mL of extracts or standard solution of Catechin (150, 200, 300, 400 and 500 mg/lit) was added to 25 mL volumetric flask. To this flask, 10 mL of distilled water was added. Further 0.75 mL of 5 % sodium nitrite (NaNO2) was added. To this solution mixture, 0.75 mL of 10 % aluminium chloride (AlCl₃) was added after 5 minutes. At the 6th min, 2 mL of 1 M sodium hydroxide (NaOH) was added and the total volume was made up to 25 mL with distilled water.

Similarly, the reagent blank was prepared by omitting the samples to be analysed. The solutions were mixed well, and the absorbance was measured against prepared reagent blank at 510 nm in UV-VIS spectrophotometer. The flavonoid content expressed as mg Catechin equivalents/gr. weight of extract.

The total flavoniodal content was calculated by using the formula



3. RESULTS AND DISCUSSION

3.1 In-Vitro Antioxidant Activity

The DPPH assay establishes the capability of extracts (Ethyl acetate, Methanol and aq. methanol) to reduce DPPH radical to the hydrazine by changing the unpaired electrons to the paired electrons. In the present study, dose- dependent inhibition of DPPH radical indicates that the extracts caused the reduction of DPPH radical. Percentage scavenging of DPPH radical by the extracts (Ethyl acetate, Methanol and aq. Methanol extracts) examined at different concentrations is depicted in Table-1. Fig-1 illustrates the effect of Ethyl acetate, Methanol and aq. Methanol extracts on DPPH radicals.

Table 1: Effect of Ethyl acetate, Methanol, and 30 % aq. methanol extracts of T. stans leaf and stem onDPPH radicals.

			% Inhibition of DPPH by anti- oxidant in the extracts					
c Concentr	Ascorbic	Leaf Extra	Leaf Extracts			Stem Extracts		
S. No	ation (µg/ml)	Acid	Leaf Ethyl acetate	Leaf Methanol	Leaf Aq. Methanol	Stem Ethyl acetate	Stem Methanol	Stem Aq.Methanol
1	5	38.23	0.18	8.35	0.90	1.81	18.79	3.09
2	10	58.56	6.81	14.45	11.36	2.81	34.16	3.09
3	15	74.58	8.90	27.81	18.62	10.31	39.68	7.93
4	20	93.92	10.89	34.50	26.64	11.71	49.03	11.69
5	25	95.02	18.52	39.59	35.42	20.25	57.22	14.95
6	50	97.23	39.87	76.52	60.56	37.42	60.31	37.67

International Journal of Pharmacy and Biological Sciences

R.Prathibha* et al 964



The Anti-oxidant activity of the *T.stans* leaf and stem extracts using DPPH method was observed to be more in stem methanolic extract when compared to other extracts. The order of Anti-oxidant activity of plant extracts by DPPH assay method is-

Stem Methanol> Leaf Methanol > Leaf Aq.Methanol > Stem Ethyl acetate > Leaf Ethylacetate > Stem Aq.Methanol





Figure 1: Graph of free radical scavenging activity (Anti-oxidant activity) of *T. stans* leaf and stem ethyl acetate, methanol and aqueous methanolic extracts on DPPH free radicals.


3.2. Total Phenolic Content

The total phenolic content of *T. stans* leaf and stem extracts were estimated by Folin-Ciocalteu's assay method. The results of the phenolic content of *T. stans* leaf and stem extracts (Hexane, Ethyl acetate, Methanol and 30% aq. Methanolic extracts) equivalent to 7.77,

54.61, 60.24, 58.34, 7.27, 54.12, 56.28 and 41.56 mg Gallic acid equivalents/gram of extract respectively were detected and the same were depicted in Table-3. The Standard graph of gallic acid is depicted in Table 2, Fig-2

Table 2: Absorbance values of Standard Gallic Acid					
	Concentration (µg/ml)	Absorbance			
	0	0			
	50	0.258			
	100	0.519			
	150	0.761			
	200	1.034			
	250	1.31			





Int J Pharm Biol Sci.

S. No	Extract	Unknown	mg/gm gallic acid	
	Extract	Concentration (µg/ml)	equivalents	
1	Leaf Hexane	77.78	7.77	
2	Leaf Ethyl acetate	546.19	54.61	
3	Leaf Methanol	602.44	60.24	
4	Leaf Aq.Methanol	583.43	58.34	
5	Stem Hexane	72.79	7.27	
6	Stem Ethyl acetate	541.20	54.12	
7	Stem Methanol	562.89	56.28	
8	Stem Aq.Methanol	415.65	41.56	

Table 3: The phenolic content of Hexane, Ethyl acetate, Methanol and 30%aq. methanol extracts of T. stans leaf and stem in terms of mg/gm gallic acid equivalents.

Based on the results, it is observed that in *T. Stans* the leaf Methanolic extract contains more phenolic content (60.24 mg GAE/ gr. Ext.) compared to other extracts. The order of Phenolic content of plant *T. stans* leaf and stem extracts-


Leaf Methanol > Leaf Aq.Methanol > Stem Methanol > Leaf Ethyl acetate > Stem Ethyl acetate > Stem Aq.Methanol > Leaf Hexane > Stem Hexane


3.3. Total Flavonoid Content

The total flavonoid content using quercetin and catechin as standards:

The total flavonoid content of *T. stans* leaf and stem extracts were estimated by aluminum chloride

colorimetric assay method using Quercetin and Catechin as standards. The leaf and stem extracts such as Hexane, Ethyl acetate, Methanol and aq. Methanolic extracts showed the results equivalent to 62.92, 189.51, 324.80, 271.01, 109.05, 221.98, 241.17 and 124.91 mg/100-gram equivalents respectively of Quercetin and equivalent to 26.69, 65.87, 123.90, 127.18, 44.38, 69.47, 95.53 and 42.90 mg/100gram equivalents respectively of catechin. The results are depicted in table- 5. The standard graphs of quercetin and catechin were depicted in Table-4, Fig-3.







The Total Flavonoid content of extracts using quercetin as standard is observed to be more in leaf methanol extract (324.80 mg QE/gr. Ext.) when compared with the other extracts. The order of Flavonoid content of plant extracts using quercetin as the standard -Leaf Methanol > Leaf Aq. Methanol > Stem Methanol > Stem Ethyl acetate > Leaf Ethyl acetate > Stem Aq.

Methanol > Stem Hexane > Leaf Hexane

The Total Flavonoid content of *T. stans* leaf and stem extract using catechin as standard was observed to be more in leaf aq. methanol extract (127.18 mg CE/gr. Ext.) when compared with the other extracts. The Order of Flavonoid content of plant extracts using catechin as standard-

Leaf Aq. Methanol > Leaf Methanol > Stem Methanol > Stem Ethyl acetate > Leaf Ethyl acetate > Stem Hexane > Stem Aq. Methanol > Leaf Hexane

Table 5: The total flavonoidal content of Hexane, Ethyl acetate, Methanol, and 30% Aq. Methanol extracts
of Tecoma stans leaf and stem in terms of mg/gm quercetin and catechin equivalents.

	Extract	Catechin		Quercetin		
Plant Part		Unknown Concentration (μg/ml)	Flavonoid Content mg Catechin Equivalent/gr. Extract	Unknown Concentration (μg/ml)	Flavonoid Content mg Quercetin Equivalent/gr. Extract	
	Hexane	266.96	26.69	629.27	62.92	
T. stans	Ethyl					
Leaf	acetate	658.75	65.87	1895.19	189.51	
Lear	Methanol	1239.04	123.90	3248.01	324.80	
	Aq. MeOH	1271.85	127.18	2710.16	271.01	
	Hexane	443.88	44.38	1090.05	109.05	
T . 4	Ethyl					
<i>T. stans</i> Stem	acetate	694.78	69.47	2219.87	221.98	
	Methanol	955.33	95.53	2411.72	241.17	
	Aq. MeOH	429.08	42.90	1249.11	124.91	

4. SUMMARY AND CONCLUSION

The extraction of T. stans leaf and stem using different solvents such as hexane, ethyl acetate, methanol, and aq. methanol was carried out. The antioxidant activity performed using DPPH method showed the stem methanolic extract as the most active one exhibiting free radical scavenging property with the percentage value of 57.22 µg/ml at 25 µg/mL concentration. The stem methanolic extract contains most effective antioxidant phytochemicals. Similarly, the total phenolic content was performed by Folin-Ciocalteu's assay method using Gallic acid as a standard. The leaf methanolic extract contained more amount of phenolic content (60.24 mg GAE/gr. Ext.). Likewise, the total flavonoidal content was performed by aluminium chloride colorimetric assay method using Quercetin and Catechin as the standards. The leaf methanolic extract showed a high amount of flavonoidal content in terms of quercetin equivalents (324.80 mg QE/gr. Ext.) whereas the leaf aq.methanol extract showed a high

amount of flavonoid content (127.18 mg CE/ gr. Ext) in terms of catechin equivalents.

ACKNOWLEDGEMENT

Authors thank Dr.G.Krishna Mohan, Professor, Centre for Pharmaceutical Sciences, Institute of Science and Technology, JNTU, Hyderabad for his support and encouragement.

5. **REFERENCES**:

- Pelton J (1964). A survey of the ecology of *Tecoma stans*. Butler University Botanical Studies, 14:53-88.
- Bailey, L.H (1941). The standard cyclopedia of horticulture, Vol. 3, MacMillan Company, New York: 2423-3639.
- Archana Singh, Nagori B.P, Kumkum Mathur (2013). *Tecoma stan*: An Important Medicinal Plant. International Journal of Pharmaceutical Erudition 3(2): 13-21.
- 4) Aguilar-Santamaría L, Ramirez G, Nicasio P, Alegria-Reyes C, Herrera-Arellano A (2009). Antidiabetic activities



of *Tecoma stans* (L.) Juss. ex Kunth. Journal of Ethnopharmacology 124 (2) : 284-288.

- Binutu OA, Lajubutu BA (1994). Antimicrobial potentials of some plant species of the Bignoniaceae family. African Journal of Medicine and Medical Sciences 23(3):269-273.
- M. Thirumal, G. Kishore, R. Prithika, Sampa Das and G. Nithya (2012). *In vitro* anticancer activity of *Tecoma stans* (L) ethanolic leaf extract on human breast cancer cell line (MCF-7). International journal of pharmaceutical, chemical and biological sciences 2(4): 488-493.
- 7) Rasika C. Torane, Sangita M. Lavate, Ravindra B. Jadhav, Gayatri S. Kamble And Nirmala R. Deshpande (2011). Evaluation of Phenol and Flavonoid Content from Aerial Parts of *Tecoma Stans*. International Journal of Pharmacy and Pharmaceutical Sciences 3 (4): 126-127.
- Raju S., Kavimani S, Uma Maheshwara rao V, Sreeramulu Reddy (2011). *Tecoma stans* (L.) Juss. ex Kunth (Bignoniaceae): Ethnobotany, Phytochemistry and Pharmacology. Journal of Pharmaceutical and Biomedical Sciences, 8(8):1-5.
- 9) Shanmukha Ittagi, Vijay Kumar Merugumolu, Ramachandra Setty Siddamsetty (2014). Cardioprotective effect of hydroalcoholic extract of Tecoma stans flowers against isoproterenol induced myocardial infarction in rats. Asian Pacific Journal of Tropical Disease 4(1): 378-384.
- Das C, Mohanty A, Sahoo DC, Dash S (2010). Evaluation of Methanolic Bark Extract of *Tecoma Stans* Linn, for Wound Healing in Albino Rats. International Journal of Pharmacy&Technology,2(3):735-742.
- Gharib Naseri MK, Asadi Moghaddam M., Bahadoram S (2007). Antispasmodic effect of *Tecoma stans* (L.) Juss leaf extract on rat ileum. Daru-journal of faculty of pharmacy 15, (3): 123.
- 12) Govindappa M, Sadananda TS, Channabasava R, Jeevitha MK, Pooja KS And Vinay B. Raghavendra, (2011). Antimicrobial, antioxidant activity and phytochemical screening of *Tecoma stans* (I.) Juss. Ex kunth. Journal of Phytology and Phytopharmacology. 3(3): 68-76.
- Marzouk M, Gamal- Eldeen A, Mohammad M, El-Sayed M, (2006). Anti-proliferative and antioxidant constituents from *Tecoma stans*, Z. Naturforsch 61: 783-91.
- 14) Jinqiu Zhu, Rene vinas, Ernest E Smith (2008). In vitro evaluation of human liver cancer cells and the potential cytotoxicity of *Tecoma stans* (Bignoniaceae) and *Brickellia cavanillesi* (Asteraceae) both single and in

Received:06.08.18, Accepted: 08.09.18, Published:01.10.2018

combination. Toxicological & Environmental Chemistry 90 (4):801 – 808.

- 15) Indra Gandhi M and Ramesh S (2010). Antifungal and haemolytic activities of organic extracts of *Tecoma stans* (Bignoniaceae). Journal of Ecobiotechnology 2(2): 26-32.
- 16) Govindappa M, Sadananda TS, Channabasava R and Vinay B. Raghavendra (2011). *In Vitro* Anti-Inflammatory, Lipoxygenase, Xanthine Oxidase and Acetycholinesterase Inhibitory Activity of *Tecoma Stans* (L.) Juss. Ex kunth, International Journal of Pharma and Bio Sciences 2(2):275-285.
- Dohnal B (1976). Investigations on some metabolites of *Tecoma stans* Juss. Callus tissue. Acta Societatis Botanicorum Poloniae 45:369-379.
- Dohnal B (1977). Investigations on some metabolites of *Tecoma stans* Juss. Callus tissue. Acta Societatis Botanicorum Poloniae 46:187-199.
- Costantino L, Raimondi L, Pirisino R, Brunetti T, Pessotto P, Giannessi F, Lins AP, Barlocco D, Antolini L, El-Abady SA (2003). Isolation and pharmacological activities of the *Tecoma stans* alkaloids. II Farmaco 58: 781-785.
- 20) Divya Sri Gurram, A. Narendra Babu, M. Sathish Kumar, V. Venkateswarlu, K. Ashok kumar (2014).
 Pharmacognostical Characteristics and Medicinal Uses of *Tecoma Stans*: A Review. Journal of Medical and Pharmaceutical Innovation 1(2) :1-4.
- 21) Darsan B Menon, and J. M. Sasikumar (2011). Antioxidant and anti-inflammatory activities of the root of *Mussaenda glabrata*. Journal of Pharmacy Research 4(10):3320-3322.
- 22) Maria Atanassova and Silviya Georgieva (2010). Comparative Polyphenol Composition and Antioxidant Capacity of The Bulgarian Plants (Dry Herbs). Electronic Journal of Environmental Agricultural and Food Chemistry 9 (9): 1514-1523.
- 23) Meetali Deori, Dipali Devi & Rajlakshmi Devi (2014). Nutrient Composition and Antioxidant Activities of Muga and Eri Silkworm Pupae. International journal of science and nature 5 (4): 636-640.
- 24) Ahlem Rebaya, Souad Igueld Belghith, Beatrice Baghdikian, Valerie Mahiou Leddet , Fathi Mabrouki, Evelyne Olivier, Jamila kalthoum Cherif, Malika Trabelsi Ayadi (2014). Total Phenolic, Total Flavonoid, Tannin Content, and Antioxidant Capacity of Halimium halimifolium (Cistaceae). Journal of Applied Pharmaceutical Science 5 (01): 052-057.

*Corresponding Author:

R.Prathibha Email: ramisettiprathibha@gmail.com