



Review of Treatment of PCOD in *Ayurved* and Allopathy

***Patil Veena Ajay¹, Khot Bhagyashree Mahadev², Kubade Nishigandha Arun³ and Gaikwad Jayant Jeevan⁴**

¹M.S. PhD, (Stree-roga Prasutitantra), Govt. Ayurvedic college, Osmanabad.

²M.S. PhD, (Stree-roga Prasutitantra), Govt. Ayurvedic college, Osmanabad.

³P.G. Scholar (Stree-roga Prasutitantra), Govt. Ayurvedic college, Osmanabad.

⁴P.G. Scholar (Stree-roga Prasutitantra), Govt. Ayurvedic college, Osmanabad.

Received: 02 Jul 2019 / Accepted: 9 Aug 2019 / Published online: 1 Oct 2019

*Corresponding Author Email: drveenapatil810@gmail.com

Abstract

The term Polycystic Ovarian Disease (PCOD) was first described by Irving Stein and Michael Leventhal as a Triad of 'Amenorrhea', 'Obesity' and 'Hirsutism' In 1935 when they observed the relation between obesity and reproductive disorders. It is hence also known as the 'Stein-Leventhal Syndrome' or 'Hyperandrogenic Anovulation' (HA) and is the most common endocrine ovarian disorder affecting approximately 2-8% women of reproductive age worldwide. Nowadays, it is also referred to as the 'Syndrome O' i.e. Over nourishment, Overproduction of insulin, ovarian confusion and Ovulatory disruption. So, PCOD is called as Polycystic Ovarian Syndrome (PCOS). PCOS is one of the most endocrinopathy affecting women. The Rotterdam 2003 criteria defines PCOS as incidence of any two of 3 key criteria namely, oligo ovulation and anovulation, hyperandrogenism and polycystic ovaries [PCO]. PCOS is found to be the most common reason for menstrual irregularities in 4-12% of women of reproductive age [12-45 yrs old]. 5-10% of women develop PCOS during their teenage or childbearing years.

Worldwide PCOS affects up to 6-7% of the population. However, the prevalence specific to the country vary extensively. In India, the incidence of PCOS/PCOD is on the hike; nearly 35% of women suffer from it. Symptoms of PCOS/PCOD are irregular, infrequent periods within 3 or 4 years of starting to menstruate light or very heavy bleeding during period weight gain, excessive hair growth to varying degrees on face, chest, and lower abdomen.

In this review paper the treatment of PCOS with *Ayurveda* and Allopathy are taken into account to compare and analyze best curable medicinal system for PCOS. Allopathy does not cure PCOS but helps in managing and controlling effects while *Ayurveda* can be considered as best cure and promising treatment with no side effects.

Keywords

PCOS/PCOD, polycystic ovaries, syndrome, follicles, cysts, menses, GnRH, homeopathy, *ayurveda*, allopathy, diabetes mellitus, anovulation, hormone.

INTRODUCTION

Polycystic Ovarian Syndrome also known as PCOS OR PCOD [Polycystic Ovarian Disorder] is a very common hormonal disorder and a leading cause of female infertility worldwide [1]. PCOS is also called as Stein-Leventhal Syndrome after two doctors who first described it in 1935. PCOS is one of the most endocrinopathy affecting women [2]. The Rotterdam 2003 criteria defines PCOS as incidence of any two of 3 key criteria namely, oligo ovulation and anovulation, hyperandrogenism and polycystic ovaries [PCO][3]. Polycystic Ovarian Syndrome (PCOS) is a condition in which women typically have many numbers of small cysts around the edge of their ovaries. Polycystic ovaries mean the ovaries containing a large number of cysts that are not bigger than 8mm and develop more follicles than normal every month. Polycystic ovary starts maturing at least twice as many follicles compared normal most of which enlarge and mature but do not release an egg. The cysts are the egg containing follicles that do not develop properly because of hormonal imbalance [4]. Some women go on to develop PCOS [Polycystic Ovarian Syndrome] which means they have other symptoms including polycystic ovaries. PCOS may be hereditary as well. Studies show that women with family history of polycystic ovaries are 50% more likely to develop PCOS. PCOS is found to be the most common reason for menstrual irregularities in 4-12% of women of reproductive age [12-45 yrs old]. 5-10% of women develop PCOS during their teenage or childbearing years. Worldwide PCOS affects up to 6-7% of the population. However, the prevalence specific to the country vary extensively [5]. In India, the incidence of PCOS/PCOD is on the hike; nearly 35% of women suffer from it [6]. Symptoms like irregular, infrequent periods within 3 or 4 years of starting to menstruate, lighter or very heavy bleeding during period, weight gain, excessive hair growth to varying degrees on face, chest, and lower abdomen. Moderate abdominal discomfort during periods, acne, and excessive skin growth on neck or in armpit also called as skin tags. Bone pain [arthralgia] and hair loss [alopecia], constipation, flatulence, and indigestion is also seen in some patients [7]. PCOS is also called as polycystic ovary disease [PCOD], Stein-Leventhal syndrome, ovarian hyperthecosis and sclerocystic ovary syndrome.

Pathogenesis

Complete understanding of pathogenesis of PCOS is still lacking due to heterogeneity of this disorder. There are most likely multiple underlying pathophysiological mechanisms. Various theories

have been proposed to explain the pathogenesis of PCOS/PCOD. They are as follows:

- a) An alteration in gonadotropin releasing hormone secretion results in increase of LH secretion. [Luteinizing hormone]
- b) An alteration in insulin secretion; leads to hyperinsulinemia and insulin resistance.
- c) Defect in androgen synthesis that leads to increase in ovarian androgen production.

LH (Luteinizing Hormone): LH hyper secretion is a main cause of infertility and miscarriage in women having PCOS/PCOD. Many theories have been introduced for the etiology of over secretion of LH by pituitary gland. These include hypothalamic dysfunction, reduced pituitary sensitivity to gonadotropin releasing hormone [GnRH] and increased pulsatility of GnRH.

Increase in LH leads to increase in androgen production by theca cells within the ovary [8].

Hyperandrogenism or Androgen excess: Hyperandrogenism is one of the primary symptoms of PCOS/PCOD. Increased or elevated level of circulating androgen is observed in 60-80% of women with PCOS [9].

Clinical features of hyperandrogenism in women with PCOS include acne, hirsutism, and androgenic alopecia [hair loss]. Insulin may modulate Gonadotropin secretion.

Hyperinsulinemia and Insulin resistance [IR]:

Hyperinsulinemia is a condition in which there is excess level of insulin circulating in the blood relative to the levels of glucose. Hyperinsulinemia can result from various metabolic diseases and condition, one of which is PCOS/PCOD [10]. Whereas insulin resistance is a pathological state or condition in which the ability of cells to respond to normal action of hormone insulin is diminished. Insulin resistance further leads to development of Type 2 diabetes mellitus. 30-40% of women affected with PCOS have impaired glucose tolerance and 10% of women to develop type 2 diabetes mellitus by the age of 40 yrs [11]. Insulin acts collaboratively with LH to enhance androgen production in the ovarian theca cells. It also decreases hepatic synthesis and sex hormone binding globulin secretion, hormone which binds testosterone in circulation therefore increasing the amount of free testosterone which is biologically available [12]. Women with PCOS/COD and hyperinsulinemia have free testosterone, but the total concentration of testosterone may be at the upper level of normal or modestly elevated.

Causes:

The main cause of PCOS/PCOD is unknown both environmental and genetic factors are implicated.

Causes of PCOS are as follows:

1. Genetic susceptibility
2. Raised levels of insulin
3. Hormonal imbalance
4. Contraceptive pills
5. Strong stimulation in adrenal in childhood
6. Obesity
7. Hereditary factors
8. Sedentary lifestyle
9. Stress
10. Diabetes
11. Insulin resistance
12. Hyperprolactinemia
13. Cushing's syndrome
14. Congenital adrenal hyperplasia [13]

Signs and symptoms:

Symptoms of PCOS vary from women to women. Some of them are as follows:

1. Infertility- by preventing ovulation.
2. Irregular, scanty, absent menses
3. Amenorrhea [30-40% of women]
4. Oligomenorrhea [85-90% of women]
5. Hirsutism [14]
6. Hyperandrogenemia
7. Acne, oily skin, dandruff
8. Depression or anxiety
9. Pelvic pain
10. Hair loss or male pattern baldness
11. Weight gain or obesity
12. Metrorrhagia [15]
13. Swollen breasts before period
14. Bleeding with uterine fibroids during menses
15. Neuralgic pain during menses
16. Hysteria
17. Itchy vagina and vulva
18. Heavy periods
19. Sleep apnea
20. Cysts on ovaries
21. Skin tags
22. High blood pressure [16]

Progression:

The condition PCOS is a very common hormonal disorder that can occur any time in women's life. Depending on when it occurs effects vary [17]. PCOS caused during adolescence and adulthood will cause reduced or no periods or menses, polycystic ovaries, obesity, and excess sex hormone levels [18]. Whereas if caused in ageing individuals it causes diabetes, high blood pressure, abnormal blood lipid i.e. cholesterol level also called as metabolic

syndrome [19]. The term "Syndrome XX" has been coined as name for PCOS [20].

Diagnosis:

There are several tests to diagnose PCOS/PCOD. Doctor follows following steps to confirm PCOS/PCOD:

a. Medical history: Menstrual periods, weight changes and other symptoms are observed.

b. Physical Examination: Measure blood pressure, Body Mass Index [BMI], and waist size, checking the areas of increased hair growth for Hirsutism [21].

c. Pelvic Examination: Examination for enlargement of ovaries or swollen by increase number of cysts.

d. Blood Test: Blood test for hormone androgen and glucose levels [22].

e. Vaginal ultrasound sonogram/sonography: Use of sound waves to take picture of pelvic area for ovarian cysts and for checking the endometrium thickness of womb.

Treatment:

PCOS/PCOD has received scant attention in the social science literature. The approach to manage PCOS, first line of treatment is by *Ayurveda*, homeopathy and or allopathy

Ayurveda:

Ayurveda classifies PCOS as a disorder of *kapha dosha*. *Vata* is responsible for movement of follicle during ovarian cycle the rupture of the ovarian wall release the matured ovum to the movement of fimbriae the finger like projections that guide the ovum into fallopian tubes and movement of ovum towards the uterus.

PCOS is due to *kapha* blocking *vata* and *pitta*. Hence *granthiadhara* [cystic swelling], *arbuda* [glandular swelling] [tumour formation]. *Apanavayu dushti* or *margavarodhjanya* [obstructed channels and transformation process is suppressed]. *Apanavayu* in *artavavahasrota* becomes *sanga* or *stagnant* due to excessive *kapha* and *ama* accumulation blocks the channel impeding the flow of *vata* in the ovarian cycle. As *vata* is blocked, *pitta* is also blocked as well, *pitta* in order to act as the intelligence behind transformation needs the movement of *vatta* in order for its energy to have potential. The accumulated *kapha* is expressed information of cyst in the ovary as it takes on heavy white sticky quality expressing *kapha* and *ama*. Menstrual problems manifest due to aggravation of all *tridoshas* but mainly due to *apanavayu*. Therefore, it should be treated before *pitta* and *kapha* – *Astanga Hridaya*. Medicines like *daruharidra* roots and Line of treatment in which *Kanchnar guggul*, *Varunadi kwath* with bide laxative like *karanj* added to these

hypoglycemic drugs. In *panchkarma* at first *vaman*, *virechan*, *bastikriya*, *nasya* are induced. Drug or generic preparations like *ashwagandhaarishta*, *ashokarishta*, *kumariasava*, M2 Tone, *chandraprabha vati*, *abhrak bhasma*, *raupya bhasma*, *phal ghrita* with moderate exercise like *pranayama*, *simhasana*, *vjrasana*, *shalabhasana*, *sarvangasana* is also helpful. The treatment is long term in nature apart from being cost effective with no side effects.

Low-fat milk is recommended. Boil milk before drinking. Drink it warm. This makes it easier to digest. Do not take milk with a full meal or with sour or salty foods. Add a pinch of turmeric or ginger to the milk before boiling it. This helps to reduce the *kapha* increasing qualities of the milk.

Lighter fruits, such as apples, oranges, grapefruit, pineapple and papaya and pears, are recommended. Sweeteners: Honey is excellent for reducing *kapha*. Reduce the intake of sugar products, which increase *kapha*.

Beans are recommended. Nuts should be avoided.

Grains such as barley, chickpea and millet are recommended. Do not take too much wheat or rice, as they increase *kapha*.

Spices are recommended only in moderate quantities.

Vegetables such as potatoes and sweet potatoes should be avoided as they can cause weight gain.

Follow a lifestyle that will not aggravate any *dosha*.

Negative feelings like stress and tension should be avoided. Do not undertake physical or mental work beyond your capacity.

Limit the use of contraceptives.

Follow an active lifestyle, yet the exercises should be according to one's capacity. Both, excessive fasting and overeating are harmful.

Smoking, drinking alcohol and using narcotics aggravates menstrual disorders.

Allopathy:

Allopathy treatment depends on basis of hormonal imbalance and the major treatment induces HRT [Hormone Replacement Therapy] where ensuing hormones are administered after evaluation of the patient.

Apart from that there is growing misconception that PCOS occurs due to malfunction of the insulin hormone and hence diabetic drugs like Metformin a oldest insulin sensitizer may also be prescribed.

Surgery is also exercised where the ovary is pierced or perforated, or the cystic tissue is destroyed for better hormonal gain. In taking hormones it further depletes endocrinal system and puts them in sleep.

The stuff which has to be naturally released by the body once received with artificial means put the body in a lazy mode. The HRT treatment may give good results but is not recommended in long run as it causes other problems.

The drugs commonly used in allopathy for treatment in women with PCOS are metformin which can lower TSH levels and hypothyroidism.

PCOS is strongly associated with preeclampsia, premature birth, more than double risk of GDM and birth of large for gestational age infants.

As a part of treatment sometimes a drug called Clomiphene combined with low doses of Dexamethasone a steroid which suppresses androgen production from adrenal glands.

An allopathic medicine doesn't have cure, but their medicines to manage PCOS/PCOD leads to numerous side effects. If the intake of medicines is stopped PCOS/PCOD reoccurs. Root cause of the problem should be treated, and allopathic medicine fails to do that. Allopathic PCOS/PCOD treatment are not without fair share of side effects including bloating, pelvic pain, night sweat or vasomotor symptoms, blurred vision, pale yellow eyes and skin, malnutrition, heavy menstrual period or bleeding between periods.

DISCUSSION

As PCOS/PCOD is multifaceted problem with reproductive endocrine and metabolic dysfunction. PCOS is characterized by infertility, ovarian dysfunction, hyperandrogenism, insulin resistance and chronic anovulation while major metabolic consequences include obesity, type II diabetes and cardiovascular disease affecting 5-10 % of female population of developed countries. Symptoms like irregular, infrequent periods within 3 or 4 years of starting to menstruate light or very heavy bleeding during period weight gain, excessive hair growth to varying degrees on face, chest, and lower abdomen. Moderate abdominal discomfort during periods, acne, and excessive skin growth on neck or in armpit also called as skin tags. Bone pain [arthralgia] and hair loss [alopecia], constipation, flatulence, and indigestion are also common. The treatment of PCOS *ayurveda* and allopathy is taken into account to compare and analyze best curable medicinal system for PCOS. The treatment generally focuses on management of main concerns such as infertility, acne or obesity. The comparison of treatment in *Ayurveda* and Allopathy system is considered to find out the system most effective in treating PCOD/PCOS. A conventional treatment system varies or differs as per the symptoms and also

has some side effects. Comparing allopathy and *Ayurveda*, *Ayurveda* shows best curable effects compared to allopathy as allopathy only aims at managing and controlling diseases.

CONCLUSION

PCOS is an increasing public health problem which is very common and leading cause of infertility in women. Polycystic ovarian syndrome or PCOS is a condition in which a women's level of sex hormone like estrogen and progesterone are imbalanced. Symptoms like irregular, infrequent periods within 3 or 4 years of starting to menstruate light or very heavy bleeding during period weight gain, excessive hair growth to varying degrees on face, chest, and lower abdomen. Moderate abdominal discomfort during periods, acne, and excessive skin growth on neck or in armpit also called as skin tags along with Bone pain [arthralgia], hair loss [alopecia], constipation, flatulence, and indigestion. The treatment of PCOS with *Ayurveda* and allopathy is taken into account to compare and analyze best curable medicinal system for PCOS. As allopathy is a method of treating disease with remedies/medications that produce effects different from those caused by the disease. Allopathy does not cure PCOS but helps in managing and controlling effects while *Ayurveda* can be considered as best cure and promising treatment with no side effects.

REFERENCES

- Stein IF, Leventhal MN. Amenorrhoea associated with bilateral Polycystic Ovaries. American Journal of Obstetrics and Gynaecology 1935; 29:181.
- Knochenhauer ES et al. Prevalence of Polycystic Ovarian Syndrome. Journal of Clinical Endocrinology and Metabolism 1998; 83(9):3068-3082.
- Rotterdam ESHRE/ASRM- Sponsored PCOS Consensus Workshop Group. Revised 2003 consensus on diagnostic criteria and long-term health risks related to polycystic ovary syndrome. Fertil, Steril. 2004; 81:19-25.
- Knochenhauer ES, Key TJ, Kahser- Miller M, Waggoner W, Boots LR, Azziz R. Prevalence of Polycystic Ovarian Syndrome in unselected black and white women of the Southeastern United States: a prospective study. J Clin Endocrinol Metab 1998; 83:3078-3082.
- Farah L, Lazeby AJ, Boots LR, Azziz R. Prevalence of Polycystic Ovarian Syndrome in women seeking treatment from community electrologists. Alabama Professional Electrology Association Study Group. J Reprod Med 1999; 44:870-874.
- Lindholm A, Andersson L, Eliasson M, Bixo M, Sundstrom-Poromaa I. Prevalence of Symptoms associated with polycystic ovarian syndrome. Int J Gynaecol Obstet. 2008 July; 102(1):39-43.
- Miller LG, Murray WJ. Herbal medicinals: a clinician's guide. Routledge; 1998. p.326.
- Ehrmann DA. Medical progress: Polycystic ovary syndrome. J Engl J Med 2005; 352:1223-1236.
- Tsilchorozidou T, Overton C, Conway G S. The pathophysiology of polycystic ovary syndrome. Clin Endocrinol (Oxf) 2004; 60:1-17.
- Legro RS et al. Prevalence and predictors of risk for type 2 diabetes mellitus and impaired glucose tolerance in polycystic ovary syndrome: a prospective, controlled study in 254 affected women. J Clin Endocrinol Metab 1999; 84(1):165-69.
- Carmina E. Diagnosing polycystic ovary syndrome in women who menstruate regularly. Contemp Obstet Gynecol 2003; 53-64.
- Orio Jr F, Matarese G, Di Biase S, Palomba S, Labella D, Sanna V, Savastano S, Zullo F, Colao A, Lombardi G 2003 Exon 6 & 2 peroxisome proliferator activated receptor-gamma polymorphisms in polycystic ovary syndrome. J Clin Endocrinol Metab 88:5887-5892 Abstract, Medline.
- Chang WY, Knochenhauer ES, Bartolucci AA, Azziz R 2005 Phenotypic spectrum of polycystic ovary syndrome: clinical and biochemical of the major clinical subgroups. Fertil Steril 83:1717-1723 Cross Ref, Medline.
- Hahn S, Tan, Elsenbruch S, Quadbeck B, Hermann BL, Mann K, Janssen OE 2005 clinical and biochemical characterization of women with Polycystic ovary syndrome in North Rhine-Westphalia. Horm Metab Res 37:438-444 Cross Ref, Medline.
- Conway GS, Honour JW, Jacobs HS 1989 Heterogeneity of polycystic ovary syndrome: clinical, endocrine and ultrasound features in 556 patients Clin Endocrinol (Oxf) 30:459-470.
- Legro RS, Myer SER, Barnhart HX, Carr BR, Carson SA, Diamond MP, Karr BA, Schlaff WO, Coutifaris C, McGovern PG, Cataldo NA, Steinkampf MP, Nestler J E, Gosman G, Giudice LC, Lepper PC The pregnancy in polycystic ovary syndrome in unselected black and white women of the southeastern United States: a prospective study. J Clin Endocrinol Metab 83:3078-3082.
- Dunne N, Slater W. The Natural Diet Solution for PCOS and Infertility: How to Manage Polycystic Ovary Syndrome Naturally. Natural Solutions for PCOS; 2006.
- Apridonidze T et al. Prevalence and Characteristics of the Metabolic Syndrome in Women with Polycystic Ovary Syndrome. J Clin Endocrinol Metab 2005; 90(4): 1929- 35.
- Ehrmann DA. Polycystic ovary syndrome. N Engl J Med 2005; 352:1223-1236.
- El Sheikh M, Caroline M. Polycystic Ovary Syndrome. Oxford University Press; 2008. 21.
- Goldzieher JW, Green JA. The Polycystic ovary: clinical and histologic features. J Clin Endocrinol Metab 1962; 22:325.
- Sinha U, Sinharay K, Saha S, Long kumar T A, Baul, SN, Pal SK, Thyroid disorders in polycystic ovarian syndrome subjects. A tertiary hospital based cross-



sectional study from eastern india. Indian J Endocrinal Metab.2013;17(2):304-309
23. Roos N, Kieler H, Sahlin L, Ekman Ordeberg G, Falconer H, Stephansson O Risk of adverse pregnancy outcomes

in women with PCOS. Population based cohort study.BMJ.2011;343;d6309