



MEDICINAL ATTRIBUTES OF FAMILY RUBIACEAE

Sirigiri Chandra Kala*

Department of Botany and Microbiology, Acharya Nagarjuna University, Nagarjuna Nagar, Guntur 522510, Andhra Pradesh, India.

*Corresponding Author Email: kala.siri2008@gmail.com

ABSTRACT

Large number of medicinal plants has been extensively screened for biologically active molecules with an intension of finding new drugs for treating different diseases. A number of medicinal plants have been evaluated for their healing potentials most of them have shown their protective effects against several diseases. Drug discovery from medicinal plants comprise a multifaceted approach of combining Botanical, Phytochemical, Biological, and Molecular techniques. The Present study was aimed to know the importance of phytochemical, antimicrobial and pharmacological activities of Rubiaceae species, so noticed that drug discovery dependsup on the preliminary and basic analysis.

KEY WORDS

Rubiaceae, Phytochemical, Antimicrobial, Pharmacological.

INTRODUCTION

Medicinal plants have been used in traditional health care systems since prehistoric times and are still the most important health care source for the vast majority of the population around the world. Nature itself has a source of medicinal agents for thousands of years and an outstanding number of modern drugs have been isolated from natural sources, many based on their use in traditional system of medicines. [1] However, recent evidences from the pharmaceutical companies' shows that for some complex diseases, natural products still represent an extremely valuable source for the production of new chemical entities^[2]. Medicinal plants play a vital role for the development of new drugs. During 1950-1970 approximately 100 plants based drugs were introduced in USA including deserpidine, reseinnamine, vinblastine vincristne which are derived from higher plants.

In order to find new sources of plant drugs, number of plants has been screened for various biological activities in various search institutions. In India, about 3,000 plants parts from 2764 plants species have been screened for their pharmaceutical and chemotherapeutic properties. The Central Drug Research Institute (CDRI), Lucknow, India has screened over 3,800 plants for a wide range of biological activities in the past 25 years. A vast wealth of medicinal plant sources is still under utilization for curing a number of diseases [3]. Recently several authors have made pharmacological studies on different plant parts [4]

PHYTOCHEMICALS

Rubiaceae of various flowering plants called the madder family, bedstraw family or coffee family. Rubiaceae family is a large family of 630 genera and about 1300 species found worldwide, especially in tropical and warm regions. Many

International Journal of Pharmacy and Biological Sciences (e-ISSN: 2230-7605)



Available Online through www.ijpbs.com (or) www.ijpbsonline.com

Rubiaceae family plants exhibited antimalarial, antimicrobial, antihypertension, antidiabetic, antioxidant, and anti-inflammatory activities. Bioactive compounds including indole alkaloids, terpenoids and anthraquinones have been isolated from these plants. Various natural products occur in Rubiaceae plants. Extensive phytochemical investigation has been realized regarding the natural occurrence of terpenoids, anthraquinones and indole alkaloids in the family. The occurrence of alkaloids in some Rubiaceae members is well documented. Phytochemical analysis of *Canthium horridum* was reported [5]. Canthium multiflorum extracts revealed the presence of several chemical compounds such as alkaloids, terpens and tannins. [6] Chemical constituents of the stems of Canthium simile was reported [7].

Canthium mannii plant bark extracts comprise nematicidal activity and valuable secondary metabolites such as alkaloids and saponins^[8]. In species, family phytochemical Rubiaceae screening has shown that in Rytigynia nigerica some bioactive compounds such as tannins, saponins, reducing compounds, steroids, and flavonoids are present. In Rytigynia umbellulata, alkaloids, saponnins, reducing tannins, compounds, and flavonoids were present. [9] Borreria and Spermacoce species has alkaloids, iridoids, flavonoids and terpenoids^[10]. Canthium multiflorum leaves exhibits high value phytochemicals like Saponins, Tannins, Flavonids, Alkaloids, Proanthocyanidins, Anthracenosides, Coumarins, Terpenoids, Sterols and Carotenoids.

Antimicrobial activity

The antimicrobial nature of the plants has been attributed to the wide variety of compounds they synthesized. The screening of bioactive compounds has always been great interest to scientist liking for new source of drugs useful in the treatment of infectious diseases. *Canthium multiflorum* showed potent antimicrobial activity

against Escherichia coli, Enterococcus faecalis, Bacillius cereus, Proteus misrabilis, Staphylococcus aureus, Staphylococcus camorum, Shigella dysentrica, Salmonella enterica, Pseudomonas aeruginosa, and Staphylococcus pyogens of various bacteria tested for antibacterial activity. Asaseet al., 2008^[11]found more inhibition of both Gram positive and Gram negative bacteria by the acetone extract of Mitragyna inermis. Adomi, 2008^[12] screened and observed high zone of inhibition using the aqueous extract of Morindalucida. The wild plants ethanolic extract of Canthium coromandelicum shows a broad spectrum of antimicrobial activity against Salmonella typhi and antifungal activity showed against Candida albicans.[13]

Pharmacological activities

Borreria and Spermacoce species of Rubiaceae as well as their isolated compounds possess diverse biological activities, including anti-inflammatory, antitumor, antimicrobial, larvicidal, antioxidant, gastrointestinal, anti-ulcer, and hepato protective, with alkaloids and iridoids as the major active principles.^[14]

Rubiaceae species were a valuable source of new secondary metabolites for medical purposes. Reports for biological activity of Rubiaceae species are numerous, but phytochemical investigations have been conducted only on a few species such latifolia, Nauclea pobequinii, Nauclea Mitragyna inermis, Pterocarpusbussei. Crude extracts of these plants have been found to have antibacterial activity. Indeed, antimicrobial property of Rubiaceae may become an useful tool in treating opportunistic infection.

Cytotoxicity

About 3000 plants has anticancer properties are subsequently used as potent anticancer drugs. Based on the previous reports, it is realized that the Rubiaceae species such as *Morinda lucida* and *Nauclea latifolia* has showed potent cytotoxic activities. [15]

IJPBS | Volume 5 | Issue 2 | APR-JUN | 2015 | 179-181

CONCLUSION

However, recent evidences from the pharmaceutical companies' shows some complex diseases, natural products still represent an extremely valuable source for the production of new chemical entities. Phyto pharmacological screening of medicinal plants and their extracts will reveal their presence of valuable compounds and provide insight into new ways of treatment with new drugs.

REFERENCES

- Ruban P, Gajalakshmi K (2012), In vitro antibacterial activity of Hibiscus rosa— sinensis flower extract against human pathogens. Asian pacific journal of tropical biomedicine, 399-403.
- Calixto Joao B. (2005), Twenty-five years of researches on medicinal plants in Latin America A personal view, Journal of Ethno pharmacology, 100: 131-134.
- Sirigiri Chandra Kala. A Review on phytochemical analysis by using callus extracts of important medicinal plants. Indo American Journal of Pharmaceutical Research, 2014:4(07). 3236-3248.
- Johnson, M, Wesely, E.G., Kavitha, M.S. and Uma V. (2011). Antibacterial activity of leaves and inter nodal callus extracts of *Mentha arvensis* L. Asian Pacific J. Trop. Med., 196-200.
- Yang, W., de Oliveira A.C, Goodwin, I., Schertz, K., Bennetzenm J.L. (1996) Comparison of DNA marker technologies in characterizing plant genome diversity: variability in Chinese sorghums. CropSci 36:1669–1676.
- 6) Akomo, E.F.O., Zongo, C., Karou, S.D., Obame, L.C., Savadogo, A., Atteke, C. and Traore, A.S.(2009). *In-vitro* Anti plasmodial and Antibacterial Activities of *Canthium multiflorum* Schtum and Thonn (Rubiacea) Extracts. Pakistan Jour. Biol. Sci, 12(12): 919-923.
- 7) Wang An-wei, Chen Guang-ying, Yin Wen-qing, Han Shan-ri, Zhang yong-qiang. (2007). Chemical

- constituents from the stems of *Canthium simile* Merr. & Chur. Chemistry and Industry of Forest Products 02-05.
- 8) Wabo Pone J., MpoameMbida (2011). Acute and Sub acute toxicity of ethanol extract of *Canthium mannii* Hiern stem bark of *Mus musculus*. Indian J. Experimental Biol., 49: 146-150.
- 9) Glory O. Ajayi, Akeem B. Kadiri, Mabel E. EgbediandOluwakemi O. Oyeyemi (2011). Pharmacognostic study of two medicinal species of *Rytigynia* (Rubiaceae) from Nigeria. Phytologiabalcanica17(3): 355-359.
- Conserva, L.M., Ferreira, J.C. Jr.(2012). Borreria and Spermacoce species (Rubiaceae): A review of their ethno medicinal properties, chemical constituents and biological activities. Pharmacogn Rev. Jan. 6(11):46-55. doi: 10.4103/0973-7847.95866.
- 11) Asase, A., Kokubun, R.J. Grayer, G. Kite, M.S.J. Simmonds, A.A.O. Yeboah and Odamtten, G.T. (2008). Chemical constituents and antimicrobial activity of medicinal plants from Ghana: Cassia sieberiana, Haemato staphisbateri Mitra gynainermis and Pseudocedre lakotschyi. Phytother. Res., 22: 1013-1016.
- 12) Adomi, P.O. (2008). Screening of the leaves of three Nigerian medicinal plants for antibacterial activity. Afr. J. Biotechnol., 7: 2540-2542.
- 13) Sathish Kumar, T.Shanmugam, S.Palvannan, Bharathi Kumar, V.M. (2008). Evaluation of antioxidant properties of *Canthium parviflorum* Lam. Leaves, Natural Product Radiance, 7(2): 122-126.
- 14) Conserva, L.M., Ferreira, J.C. Jr. (2012). Borreria and Spermacoce species (Rubiaceae): A review of their ethno medicinal properties, chemical constituents and biological activities. Pharmacogn Rev. Jan. 6(11): 46-55. doi: 10.4103/0973-7847.95866.
- 15) Simplice D. Karou, TchadjoboTchacondo, Denise P. Ilboudo and Jacques Simpore, (2011). Sub-Saharan Rubiaceae: A Review of Their Traditional Uses, Phytochemistry and Biological Activities. Pakistan Journal of Biological Sciences 14: 149-169.



*Corresponding Author: kala.siri2008@gmail.com

181